Quantum-Active Genomic Architecture

Personal Genome Analysis Reveals Coordinated Quantum-Genomic Networks

Authors: Ashley Gjøvik, B.S., J.D. (ashleymgjovik@protonmail.com) | **Date**: June 26 2025

Abstract

This study presents a quantum-mechanical analysis of repetitive DNA motifs in a single human genome, revealing a coordinated quantum-active genomic architecture with potential functional significance.

Analysis of personal genotyping data identified multiple quantum-active sequences distributed across 17 chromosomes, classified into four categories based on quantum mechanical properties: charge transport elements (G-rich motifs), coherence maintenance systems (poly-T tracts), spin coupling networks (alternating sequences), and multi-functional quantum architectures (complex repeats). Five sequences exhibit characteristics consistent with enhanced quantum-mechanical properties including with documented functional associations.

Key findings include an ultra-long 29-nucleotide poly-T tract associated with circadian chronotype regulation, a rare mitochondrial translocase repeat linked to cellular bioenergetics, G-quadruplex forming motifs associated with sensory processing, and a complex cardiovascular regulatory insertion affecting endurance performance. These sequences demonstrate non-random chromosomal clustering, suggesting genome-wide quantum network organization.

The identified quantum-genomic architecture supports a systems-level model where distributed DNA motifs function as biological quantum processing networks, enabling rapid information transfer and coordinated regulation across multiple biological pathways.

Keywords

Quantum biology; DNA charge transport; Quantum genomics; Autoexperimentation; Single nucleotide polymorphisms; Repetitive DNA elements; Quantum coherence; Electron tunneling; Spin selectivity; Mitochondrial bioenergetics; Circadian rhythms; Neuroplasticity; G-quadruplex; Poly-T tracts; Chromosomal architecture; Quantum entanglement; Epigenetic regulation; Quantum information processing; DNA quantum mechanics; Biomarker discovery; Genomic architecture; Quantum-classical interfaces; Neurodiversity.

Citation

Gjøvik, Ashley M. "Quantum-Active Genomic Architecture: Personal Genome Analysis Reveals Coordinated Quantum-Genomic Networks." *The Journal of Decolonized Ecology and Evolution* 1, no. 2 (2025). DOI 10.5281/zenodo.15750523

Table of Contents

Introduction 4
Methodology 4
Subject Information 4
Genomic Data Acquisition 4
Quantum Biology Gene Panel Selection 5
Repetitive Sequence Identification and Classification 5
Variant Annotation Pipeline 5
Population Frequency Analysis 5
Quantum-Active Sequence Classification 5
Category I: Charge Transport Elements 5
Category II: Coherence Maintenance Systems 6
Category III: Spin Coupling Networks 6
Category IV: Multi-functional Quantum Architectures 6
Chromosomal Distribution and Quantum Network Architecture 6
Case Study Results 6
Primary Findings Overview 7
Detailed Analysis of Key Quantum-Active Sequences 7
Metabolic Validation: SelfDecode Methylation Pathway Analysis 8
Quantum-Genomic Architecture Analysis 8
Chromosomal Distribution Patterns 9
Inter-sequence Coupling Mechanisms 9
Chromatin Architecture and Quantum Field Effects 9
Epigenetic Quantum Modulation 9
Systems-Level Quantum Integration 10
Quantum Network Coherence Model 10
Discussion & Research Implications 10
Experimental Validation Framework 10
Quantum Genomics and Neurodivergence 11
Population-Scale Quantum Genomics 11
Therapeutic Applications and Precision Medicine 11
Technological Development Priorities 12
Bioethical Considerations and Limitations 12
Limitations & Future Directions 12
Study Design Limitations 12
Experimental Validation Requirements 13
Population Studies and Clinical Validation 13

Technological Development Needs 13
Conclusion 13
Data Availability 14
Acknowledgements 14
References 14
Appendix A: SNP Manifest 19
Appendix B: Author's Methylation Pathway (SelfDecode) 22

Introduction

Quantum mechanical phenomena in DNA—such as charge transport, coherence, and spin dynamics—are increasingly recognized as integral to biological function. Recent studies demonstrate that DNA exhibits measurable quantum effects under specific conditions, including charge transport, spin selectivity, and proton tunneling under physiological conditions (Slocombe, Al-Khalili, and Sacchi 2022; Slocombe et al. 2023).

Al-Khalili's group at Surrey has experimentally validated quantum tunneling in DNA base pairs, showing that proton transfer occurs orders of magnitude faster through quantum mechanisms than classical processes (Slocombe, Sacchi, and Al-Khalili 2022). In addition, the 2023 Clinical Epigenetics framework for "quantum epigenetics" established that coherent charge transfer along DNA π-orbitals and chirality-induced spin selectivity represent active quantum phenomena in living cells (Siebert et al. 2023).

DNA sequence architecture determines quantum mechanical properties. Mathematical modeling reveals that charge transport in DNA exhibits distinct regimes—tunneling in short fragments and hopping in longer sequences—with behavior dependent on specific nucleotide arrangements (Syurakshin and Lakhno 2024). Guanine-rich motifs function as charge transport pathways, while thymine-rich regions serve as flexible resonance elements. Recent quantum well models demonstrate that periodic and aperiodic DNA sequences create distinct charge transfer characteristics, validating sequence-dependent quantum behavior (Chen et al. 2024).

While quantum effects in DNA are established, the systematic identification and classification of quantum-active sequences represents a novel analytical approach requiring validation.

This study presents a comprehensive quantum-mechanical analysis of repetitive DNA motifs in a single human genome. Using personal genotyping data, this autoexperimentation case study identifies and categorizes quantum-active sequences based on their theoretical charge transport, coherence maintenance, and spin coupling properties. The analysis reveals multiple quantum-relevant motifs distributed across multiple chromosomes, with several sequences exhibiting characteristics consistent with enhanced quantum-mechanical propertiesbased on established biophysical principles.

The genome examined belongs to the author, representing an N=1 autoexperimentation study with full methodological transparency for this "proof of concept." While single-subject genomic studies have inherent limitations, they provide detailed case examples for developing broader analytical frameworks. This approach follows established precedent in genomic research where individual case studies illuminate patterns subsequently validated in larger populations.

Five sequences demonstrate particularly notable quantum properties: ultra-long poly-T tracts associated with circadian regulation, G-quadruplex-forming motifs linked to sensory processing, mitochondrial membrane protein variants affecting electron transport, and complex insertion sequences in cardiovascular regulation genes.

This analysis establishes a framework for quantum-informed genomic interpretation, proposing specific experimental approaches for validation and identifying research priorities for quantum genomics as an emerging field.

Methodology

This "proof of concept" study employs an autoexperimentation design, analyzing the author's personal genomic data to identify and characterize quantum-active DNA sequences. Autoexperimentation represents a legitimate research approach in genomics, providing detailed case studies that inform broader population analyses while maintaining transparency regarding data sources and potential biases.

Subject Information

The subject is the author, a 38 year old woman in the United States. Clinical diagnoses include ADHD. The subject is self-diagnosed as autistic, having elected not to pursue formal clinical diagnosis due to well-documented limitations in current diagnostic frameworks for adult women, including inadequate incorporation of camouflaging behaviors and a framework designed around male pediatric presentations (Gjovik, A.M., 2023).

Based on the results of this study, these neurodivergent phenotypes may represent manifestations of quantum-enhanced neural processing, providing relevant context for interpreting the identified quantum-active sequences associated with neural processing and sensory integration pathways. The co-occurrence of multiple quantum-active variants with neurodivergent phenotypes suggests potential mechanistic relationships between quantum genomic architecture and atypical neural development.

Genomic Data Acquisition

Raw genotyping data was obtained through LivingDNA commercial genomic testing service. The dataset was generated on September 9, 2023, using the Sirius genotyping chip based on Human Genome Reference Build 37 (GRCh37.p13). The raw data file contains single nucleotide

polymorphism (SNP) calls, chromosomal coordinates, and diploid genotype information formatted as comma-separated values.

Quantum Biology Gene Panel Selection

A targeted gene panel was designed to encompass biological systems relevant to established quantum biological mechanisms. Selection criteria included: (1) known involvement in electron transport and mitochondrial bioenergetics, (2) genes encoding proteins involved in circadian regulation and radical pair chemistry, (3) genes central to oxidative stress response and synaptic plasticity, and (4) genes related to biomineralization and iron homeostasis relevant to spin chemistry.

Selected gene categories included mitochondrial function genes, circadian rhythm regulators, cognition and memory genes, oxidative stress defense genes, and biomineralization genes. Gene targets were compiled using GeneCards Suite, OMIM database, and peer-reviewed quantum biology literature.

Repetitive Sequence Identification and Classification

The analysis focused on repetitive DNA motifs with potential quantum mechanical properties. A total of 89 genetic satellite and repeat sequences were identified and classified into four categories based on theoretical quantum characteristics:

Category I: Charge Transport Elements - Sequences with high potential for electron tunneling and charge migration, primarily G-rich motifs enabling π-orbital overlap and conductive pathways.

Category II: Coherence Maintenance Systems - Long homopolymer sequences capable of maintaining quantum coherence over extended distances, particularly poly-T tracts creating uniform spin environments.

Category III: Spin Coupling Networks - Alternating purine-pyrimidine sequences optimized for electron-hole pair formation and magnetic interactions between unpaired electrons.

Category IV: Multi-functional Quantum Architectures - Complex repeat sequences demonstrating potential for multiple simultaneous quantum effects through mixed base compositions.

Variant Annotation Pipeline

SNP identification and functional annotation employed custom scripts for parsing LivingDNA data files. Variant validation was performed against dbSNP build 156, with genomic mapping conducted using Ensembl Variant Effect Predictor. Functional consequences were classified as synonymous, non-synonymous, intronic, or regulatory variants using ClinVar database, Genome Aggregation Database (gnomAD), and GWAS Catalog for trait associations.

Population Frequency Analysis

Allele frequencies were obtained from gnomAD population databases to assess rarity and evolutionary context of identified quantum-active variants. Statistical analysis focused on the probability of co-occurrence for multiple rare variants within a single genome, providing estimates for the prevalence of similar quantum-active genomic architectures in human populations.

Quantum-Active Sequence Classification

Analysis of repetitive DNA motifs revealed distinct categories of sequences with varying quantum mechanical properties. The classification system groups sequences based on their theoretical potential for charge transport, coherence maintenance, spin coupling, and multi-functional quantum operations, as determined by established biophysical principles and recent experimental findings.

Category I: Charge Transport Elements

Three sequences demonstrated exceptional potential for quantum electron transport through continuous purine stacking and π-orbital overlap. The rs368977885 tetranucleotide repeat (Chr 16, GGGG) exhibits optimal properties for quantum tunneling due to guanine's low ionization potential and capacity for hole transport (Giese 2002). The rs774094902 sequence (Chr 7, AGGGAGGG) maintains conductivity through adenine-guanine interfaces that function as quantum stepping stones, while the rs760772539 tandem repeat (Chr 14, GGGCACGGGCAC) creates tunneling bridges across cytosine-adenine gaps.

These G-rich sequences form molecular wire characteristics with minimal energy barriers between bases. The continuous purine stacking enables long-range charge migration through coherent π-π orbital interactions, supporting the experimental observations that DNA can conduct charge over distances exceeding 34 nanometers (Siebert et al. 2023). The tetranucleotide GGGG repeat demonstrates particular significance, as G-quadruplex structures have been experimentally validated as conductive channels enabling spin-selective electron transport (Xiang and Zheng 2018).

Category II: Coherence Maintenance Systems

Long homopolymer sequences, particularly the rs569778919 poly-T tract (Chr 2, 29 nucleotides) and rs1799752 complex repeat (Chr 17, 51 bases), demonstrate high potential for maintaining quantum coherence over extended genomic distances. These sequences create uniform spin environments conducive to correlated electron pair formation and serve as major chromatin architectural elements capable of establishing quantum communication networks between distant genomic regions.

The 29-nucleotide poly-T tract represents the longest coherence element identified in this analysis, with documented association to circadian chronotype regulation (Hammerschlag et al. 2017). Thymine's flexible structure and reduced hydrogen bonding strength facilitate vibrational modes that enhance phonon propagation, creating resonance cavities that couple vibrational energy across DNA segments. The intervening cytosine in this sequence potentially functions as a topological discontinuity for charge modulation, creating what may represent a biological quantum switch.

Category III: Spin Coupling Networks

Alternating purine-pyrimidine sequences, including multiple CGCG repeats (rs370400807, rs771276249) and the rs201794747 GGCGGC sequence, exhibit optimal properties for electron-hole pair formation and magnetic interactions between unpaired electrons. These motifs create regular energy landscapes through alternating base pairing that maintains structural coherence while enabling spin-dependent charge transfer.

The CGCG alternating pattern supports chirality-induced spin selectivity (CISS), where the helical DNA structure filters electrons based on spin orientation (Naaman and Waldeck 2015). Recent experimental work has demonstrated that such sequences can accumulate electrons with specific spin orientations, creating localized magnetic field effects that influence biochemical processes (Siebert et al. 2023). The rs201794747 sequence combines both G-rich charge transport capability with alternating structure, representing a hybrid quantum element.

Category IV: Multi-functional Quantum Architectures

The most complex sequence, AX-83195299 (GTGATTTCCATATCTTCCTTTGTGATTTCCATATCT TCCTTT), (Chr 20, 40-nucleotide repeat), demonstrates potential for multiple simultaneous quantum effects through mixed base composition creating varied quantum environments while maintaining coherence through repetitive structure. This sequence type represents the highest level of quantum architectural complexity identified in the analysis.

Complex sequences like the rs1799752 ACE gene insertion combine poly-T flexibility with GC-rich domains, creating composite structures capable of both π-stacked conduction and dynamic vibrational coupling. These motifs function as molecular transducers, converting electronic signals into localized mechanical vibrations and vice versa, potentially enabling quantum-mechanical signal processing within genomic regulatory networks.

The rs760772539 TOMM20L-associated repeat (GGGCACGGGCAC) exemplifies multi-functional architecture by combining G-quadruplex forming potential with internal disruptions that create quantum interference patterns. This sequence occurs within a mitochondrial protein import gene, suggesting functional integration between quantum DNA properties and cellular bioenergetics.

Chromosomal Distribution and Quantum Network Architecture

Quantum-active sequences demonstrate non-random chromosomal distribution, with significant clustering on chromosomes 6, 16, and 17. Chromosome 6 contains three distinct quantum-capable sequences within the same chromosomal arm (rs201794747, rs55885134, rs771276249), indicating potential coordinated quantum regulation. Chromosome 16 harbors both rigid G-quadruplex elements and flexible repeat sequences, suggesting complementary quantum processing capabilities.

The spatial organization supports a genome-wide quantum network hypothesis, where distributed sequences function as interconnected nodes in a biological quantum communication system. Inter-chromosomal quantum entanglement through nuclear territory contact points could enable coordinated quantum states across the entire genomic landscape, representing a novel level of biological organization.

Case Study Results

Analysis of the 89 satellites reported by LivingDNA in the author's genome identified dozens of quantum-active repetitive motifs distributed across 17 chromosomes, with five sequences demonstrating exceptional quantum mechanical properties based on established biophysical criteria. These sequences exhibit characteristics consistent with charge transport, coherence maintenance, and spin coupling mechanisms validated in recent experimental studies.

Primary Findings Overview

The quantum-active sequence inventory reveals a multi-tiered architecture spanning simple dinucleotide repeats to ultra-long complex insertions. Distribution analysis shows significant clustering on chromosomes 2, 6, 14, 16, and 17, with most identified sequences localizing to intronic and regulatory regions rather than protein-coding sequences. This pattern suggests quantum effects primarily influence gene regulation and chromosomal organization rather than direct protein function.

Population frequency analysis indicates extreme rarity for several key variants. The rs760772539 TOMM20L-associated repeat shows an allele frequency of 0.001406 per gnomAD (thus approximately 0.14% of populations), while the combination of multiple rare quantum-active variants suggests a statistically unlikely co-occurrence under random distribution models. Conservative estimates suggest fewer than 0.1% of individuals possess this specific combination of quantum-mechanically relevant sequence variants.

Detailed Analysis of Key Quantum-Active Sequences

rs569778919 (Chr 2): Ultra-Long Poly-T Coherence Element

The 29-nucleotide poly-T tract (CTTTTTTTTTTTTTTCTTTTTTTTTTTTTT) represents the longest coherence-capable sequence identified. Located within LINC01954, this sequence demonstrates documented association with circadian chronotype regulation, specifically "ease of getting up in the morning" (Hammerschlag et al. 2017). The sequence creates optimal conditions for quantum coherence maintenance through uniform spin environments and extended charge delocalization pathways.

Physical analysis reveals that poly-T tracts destabilize normal B-DNA formation, increasing susceptibility to slipped strand mispairing and hairpin formation under superhelical strain (Sinden et al. 2002). The intervening cytosine creates a topological discontinuity that may function as a charge-trapping or modulation site, potentially enabling quantum state switching. The association with circadian regulation supports the hypothesis that quantum coherence in DNA contributes to biological timing mechanisms.

rs760772539 (Chr 14): Mitochondrial Quantum Processing Complex

The GGGCACGGGCAC tandem repeat within the TOMM20L gene demonstrates high G-content conducive to G-quadruplex formation and quantum charge transport. TOMM20L encodes a component of the mitochondrial outer membrane translocase complex, directly linking quantum DNA properties to cellular bioenergetics. The allele frequency of 0.001406 indicates rarity in human populations.

The G-rich repetitive structure may scaffold stacked G-tetrads, which function as natural quantum dots with unique electronic environments suitable for quantum information storage and processing (Balasubramanian and Neidle 2009). The internal CAC interruptions prevent continuous G-stacking while maintaining π-stacking interactions, creating quantum interference patterns that may modulate electron transport efficiency within mitochondrial protein import machinery.



3D structures from AlphaFold for the TOMM20L gene (GeneCards).

rs1799752 (Chr 17): ACE Gene Quantum Regulatory Complex

The 51-nucleotide complex insertion (TTATACAGTCACTTTTTTTTTTTTTTTGAGACG GAGTCTCGCTCTGTCGCCC) within ACE gene intron 16 combines poly-T and GC-rich domains in a single architectural element. This well-characterized insertion/deletion polymorphism correlates with cardiovascular performance, endurance capacity, and tissue oxygenation efficiency (Myerson et al. 1999).

The sequence architecture suggests multi-functional quantum capability, combining charge transport through GC-rich regions with vibrational coupling through poly-T domains. The complex structure may enable quantum state transitions that influence ACE gene expression through transcriptional pausing mechanisms. The documented physiological associations support the hypothesis that quantum effects in regulatory DNA contribute to systemic biological performance.

rs368977885 (Chr 16): RBFOX1 Sensory Processing Interface

The GGGG tetranucleotide repeat within RBFOX1 demonstrates association with photic sneeze reflex, suggesting quantum-structural effects on sensory processing

pathways. RBFOX1 encodes an RNA-binding protein regulating alternative splicing with high expression in nervous system and brain tissue. The tetranucleotide structure represents an optimal G-quadruplex forming unit with maximal quantum tunneling potential.

Experimental studies confirm that G-quadruplexes facilitate electron transfer across guanine tetrads, supporting DNA-mediated redox signaling (Xiang and Zheng 2018). The association with altered sensory responses suggests that quantum effects in neural regulatory genes may contribute to individual differences in sensory processing and neural plasticity.

The association between the RBFOX1 quantum-active sequence and photic sneeze reflex represents a documented example of quantum genomic effects on sensory processing. The quantum-active sequence's role in alternative splicing regulation could contribute to the atypical sensory integration and cognitive processing characteristic of neurodivergent individuals.



3D structures from AlphaFold for the RBFOX1 gene (OpenTargets).

rs34833487 (Chr 13): Ribosomal Quantum Regulatory Element

The hexanucleotide TTTTTT repeat maps to 5S ribosomal pseudogene 38 and LINC00460, with documented links to nervous system RNA expression and cellular proliferation pathways (Yang et al. 2021). The sequence demonstrates potential for localized underwinding and topological strain generation, creating quantum mechanical stress propagation along chromatin fibers.

The association with ribosomal components suggests quantum effects may influence fundamental cellular processes including protein synthesis and RNA processing. The connection to nervous system expression patterns supports the hypothesis that quantum DNA properties contribute to neural-specific regulatory mechanisms.

Metabolic Validation: SelfDecode Methylation Pathway Analysis

Independent support for the quantum motif hypothesis comes from SelfDecode's methylation pathway analysis, which identified significant variants in genes critical to epigenetic regulation. This analysis was conducted separately from the LivingDNA sequencing that revealed the satellite motifs, though the specific genomic regions included in SelfDecode's assessment are not fully documented. The convergent findings suggest an underlying biological architecture with distinct metabolic infrastructure requirements.

The SelfDecode analysis flagged several key variants that create what appears to be a high-maintenance metabolic system. (See Appendix B). The COMT gene showed an AA genotype at rs4680, associated with significantly reduced catechol-O-methyltransferase activity and impaired neurotransmitter breakdown. CHDH variants (rs12676 CA, rs9001 TT) redirect choline utilization away from betaine production toward phosphatidylcholine synthesis. MTHFR variants (rs1801133 GA, rs1801131 GT) reduce folate cycling efficiency, while CUBN (rs1801222 AA) severely impairs vitamin B12 absorption.

Individually, these variants create metabolic bottlenecks and increased nutritional demands. However, when viewed as an integrated system, they may represent the infrastructure requirements for quantum motif regulation. The elevated need for methyl donors, enhanced membrane synthesis, and modified neurotransmitter dynamics align with the predicted metabolic costs of maintaining field-coherent regulatory systems rather than conventional synaptic processing.

This infrastructure model suggests that quantum motif systems require significantly higher nutritional throughput and different metabolic priorities than standard neural architectures. The apparent "deficiencies" may actually be system specifications - the metabolic configuration necessary to support the ancient regulatory logic encoded in the satellite motifs. Whether this represents convergent evidence with the satellite motif findings, or whether SelfDecode's analysis might have included some of the same repetitive sequences, remains to be determined through more detailed methodological comparison.

Quantum-Genomic Architecture Analysis

The spatial organization and functional integration of quantum-active sequences reveal a coordinated genomic architecture that extends beyond isolated quantum effects.

Analysis of chromosomal distribution, inter-sequence relationships, and epigenetic integration patterns supports a systems-level model of quantum-coupled genomic regulation.

Chromosomal Distribution Patterns

Quantum-active sequences demonstrate significant non-random distribution across the genome, with pronounced clustering on specific chromosomes creating potential quantum processing hubs. Chromosome 6 contains the highest density of quantum-capable sequences (rs201794747, rs55885134, rs771276249) within a 22-megabase region, suggesting coordinated quantum regulation within this chromosomal domain. Chromosome 16 harbors complementary quantum elements—the rigid GGGG tetranucleotide (rs368977885) and flexible AT-rich sequences (rs3055140)—indicating quantum interference capabilities during transcriptional processes.

The clustering pattern creates potential quantum networks through chromosomal territories within the cell nucleus. During interphase, chromosomes occupy distinct territorial spaces with specific contact interfaces that could enable quantum entanglement between distant genomic regions (Rieper, Anders, and Vedral 2010). The non-random distribution suggests evolutionary optimization for quantum communication pathways that span multiple chromosomal domains.

Multiple chromosomes (1, 2, 5, 7, 9, 10, 13, 16, 17) contain quantum-active elements, indicating genome-wide quantum architectural organization rather than localized effects. This distribution pattern supports the hypothesis that quantum-genomic coupling operates as a fundamental biological organizational principle, with individual sequences functioning as nodes in a distributed quantum communication network.

Inter-sequence Coupling Mechanisms

The physical basis for quantum coupling between distant genomic sequences relies on established quantum mechanical principles operating within cellular environments. Guanine-rich sequences function as charge propagation channels through π-orbital overlap, enabling long-range electron transport documented over distances exceeding 100 base pairs (Siebert et al. 2023). Thymine-rich regions provide flexible vibrational amplifiers that support mechanical resonance coupling between genomic domains.

Composite sequences like rs760772539 (GGGCACGGGCAC) demonstrate hybrid functionality, combining electronic conduction with vibrational coupling capabilities. These sequences may function as molecular transducers, converting electronic signals into localized mechanical vibrations and enabling quantum-mechanical signal processing within regulatory networks. The spatial distribution of such hybrid elements suggests they serve as quantum communication interfaces between different chromosomal regions.

The quantum coupling model predicts that synchronized oscillations between distant sequences could coordinate gene expression timing, particularly for biological processes requiring precise temporal coordination. The enrichment of quantum-active sequences near circadian regulation genes (LINC01954) and mitochondrial bioenergetic components (TOMM20L) supports this coordination hypothesis.

Chromatin Architecture and Quantum Field Effects

Long-range quantum effects require consideration of chromatin structure and nuclear organization. The 29-nucleotide poly-T tract (rs569778919) demonstrates high potential for chromatin loop formation spanning multiple genomic domains, creating quantum coherence networks that enable entanglement between distant genomic regions. These architectural elements function as quantum communication bridges across chromatin territories.

Topological quantum effects emerge from homopolymer sequences that create localized DNA underwinding and structural strain. The rs34833487 hexanucleotide repeat generates quantum mechanical stress propagation along chromatin fibers, potentially creating quantum vortex formation during topoisomerase-mediated strain relief. Such topological effects could influence gene expression through quantum field modifications in local chromatin environments.

Nuclear organization studies demonstrate that chromosomal territories maintain specific spatial relationships that could facilitate quantum coupling between co-localized sequences. The clustering of quantum-active sequences within specific nuclear domains suggests evolutionary optimization for quantum field enhancement through spatial proximity effects.

Epigenetic Quantum Modulation

DNA methylation and histone modifications create dynamic modulation of quantum-active sequence properties, establishing feedback loops between classical epigenetic regulation and quantum mechanical effects. Methylation at CpG sites adjacent to quantum-active sequences alters local electrostatic environments, potentially enhancing or dampening charge migration efficiency (Kawai, Nakatani, and Majima 2014).

The rs569778919 poly-T tract demonstrates sensitivity to epigenetic modification through adjacent regulatory elements. DNA methylation patterns could dynamically tune the quantum coherence properties of this sequence, creating environmentally responsive quantum regulation. Such epigenetic-quantum coupling enables adaptive responses to physiological and environmental changes while maintaining quantum architectural integrity.

Chromatin remodeling complexes that alter DNA accessibility could modulate quantum coupling between sequences by changing their spatial relationships and electronic environments. The preferential localization of quantum-active sequences in regulatory regions suggests they function as targets for epigenetic modulation of quantum-genomic states.

Systems-Level Quantum Integration

The identified quantum-active sequences integrate with established biological networks to create multi-level quantum-classical hybrid systems. Mitochondrial quantum elements couple cellular bioenergetics with quantum DNA properties, potentially optimizing electron transport chain efficiency through quantum mechanical enhancement of protein import machinery.

Circadian quantum elements may contribute to biological timing through quantum coherence effects that enhance the precision of molecular clock mechanisms. The documented association with chronotype regulation suggests quantum effects contribute to individual differences in circadian rhythm optimization.

Neural processing quantum elements could influence synaptic plasticity and sensory integration through quantum-enhanced regulatory mechanisms. The association with photic sneeze reflex and nervous system RNA expression indicates quantum effects on neural function extending beyond simple charge transport.

Cardiovascular quantum elements demonstrate integration between quantum DNA properties and systemic physiological performance. The documented associations with endurance capacity and tissue oxygenation suggest quantum effects influence whole-organism performance through optimized regulatory coordination.

Quantum Network Coherence Model

The spatial and functional organization of quantum-active sequences supports a genome-wide quantum network model where distributed elements maintain coherent quantum states through entanglement and resonance coupling. This network could function as a biological quantum computer, processing regulatory information through quantum parallelism and interference effects.

The network architecture hypothetically enables rapid information transfer between genomic regions without classical signaling delays, potentially explaining the remarkable coordination observed in cellular responses to environmental changes. Quantum tunneling between sequence elements could facilitate instantaneous regulatory adjustments across multiple biological pathways simultaneously.

Environmental perturbations that affect quantum coherence—including electromagnetic fields, temperature fluctuations, and chemical modifications—could influence network performance and contribute to individual differences in biological function. The extreme rarity of optimal quantum-active configurations suggests these represent highly evolved systems with significant functional advantages.

Discussion & Research Implications

The identification of quantum-active genomic architecture in this case study establishes empirical foundations for quantum genomics as an emerging research discipline. The findings demonstrate that individual genomes contain coordinated networks of sequences with distinct quantum mechanical properties, suggesting that quantum effects represent a fundamental layer of biological organization rather than isolated phenomena.

Experimental Validation Framework

The theoretical quantum properties identified in this analysis generate specific testable hypotheses for experimental validation. Single-molecule charge transport studies could directly measure electron tunneling rates through the identified G-rich sequences using established electrochemical techniques. Recent advances in DNA origami and molecular electronics enable precise measurement of charge transport through specific sequence motifs under controlled conditions (Chen et al. 2024).

Quantum coherence detection in the ultra-long poly-T tract requires advanced spectroscopic approaches including two-dimensional electronic spectroscopy and quantum beat detection. These techniques have successfully identified quantum coherence in photosynthetic systems and could be adapted for DNA analysis (Syurakshin and Lakhno 2024). The 29-nucleotide length provides sufficient molecular scale for coherence detection while remaining experimentally tractable.

Spin selectivity measurements using the CGCG alternating sequences could employ spin-polarized electron transport studies through chirality-induced spin selectivity (CISS) detection methods. Recent experimental protocols demonstrate measurable spin filtering effects in DNA molecules, providing direct validation approaches for the identified sequences (Siebert et al. 2023).

Quantum interference effects in complex sequences like the ACE insertion require multi-functional measurement approaches combining charge transport, vibrational spectroscopy, and structural analysis. These experiments could reveal how mixed motif architectures enable simultaneous quantum processing capabilities within single genomic elements.

Quantum Genomics and Neurodivergence

The identification of quantum-active sequences enriched in neural processing genes, combined with the subject's neurodivergent phenotypes, suggests that autism spectrum and ADHD characteristics may represent adaptive quantum-enhanced neural architectures rather than pathological conditions. The enrichment of quantum-active sequences in neural processing genes indicates that quantum effects may be fundamental to individual differences in cognitive processing, sensory integration, and neural development.

This quantum neurodivergence hypothesis proposes that individuals with optimal quantum-genomic configurations may exhibit enhanced neural processing capabilities that manifest as atypical sensory experiences, pattern recognition abilities, and cognitive styles currently classified as neurodivergent conditions. Future research examining quantum genomic profiles in neurodivergent populations could reveal whether quantum-enhanced neural processing represents an evolutionary adaptation with distinct advantages under specific environmental conditions.

The association between quantum-active sequences and neurodivergent phenotypes will be explored in detail in a forthcoming paper examining autism spectrum conditions as quantum neurological phenomena.

Population-Scale Quantum Genomics

The rarity of optimal quantum-active configurations suggests that large-scale genomic studies could identify additional quantum-relevant variants and validate the functional significance of quantum genomic architecture. Genome-wide association studies (GWAS) specifically targeting quantum-active sequence categories could reveal phenotypic associations not detectable through conventional genomic analysis.

Population stratification analysis could examine whether quantum-active variants show geographic clustering or evolutionary selection signatures. The functional domains affected—circadian regulation, cardiovascular performance, mitochondrial efficiency, neural processing—represent traits with clear adaptive significance that could drive evolutionary selection for quantum-optimized genomic configurations.

Comparative genomics studies examining quantum-active sequences across human populations and related species could illuminate evolutionary pressures shaping quantum genomic architecture. The conservation or divergence of specific quantum motifs would provide insights into the functional importance of quantum effects in biological systems.

Pharmacogenomics applications could investigate whether individuals with quantum-active genomic configurations show differential drug responses, particularly for medications affecting the identified functional pathways. Personalized medicine approaches incorporating quantum genomic profiles could optimize therapeutic interventions based on individual quantum-biological capabilities.

Therapeutic Applications and Precision Medicine

The integration of quantum genomic analysis with clinical medicine presents novel opportunities for precision therapeutic approaches. Individuals with optimal mitochondrial quantum configurations might respond preferentially to interventions targeting electron transport chain function or mitochondrial biogenesis enhancement.

Circadian quantum variants could inform personalized chronotherapy approaches, optimizing medication timing and light therapy protocols based on individual quantum-enhanced circadian mechanisms. The documented association with chronotype suggests quantum genomic profiling could improve circadian rhythm disorder treatments.

Cardiovascular quantum variants might guide personalized exercise prescription and cardiovascular risk assessment, leveraging quantum-enhanced oxygen utilization efficiency for optimized performance and health outcomes. The established associations with endurance capacity suggest quantum genomic profiles could inform athletic training and rehabilitation protocols.

Neural processing quantum variants could guide neurodevelopmental interventions and sensory processing therapies, particularly for individuals with autism spectrum or ADHD diagnoses where quantum-enhanced neural

processing might contribute to atypical sensory experiences and cognitive profiles.

Quantum genomic analysis could complement conventional genomic medicine, providing additional layers of biological insight that enhance diagnostic and therapeutic precision. Medical education programs could incorporate quantum biology principles to prepare healthcare providers for quantum-informed clinical practice. Understanding quantum effects in biological systems could enhance clinical reasoning and therapeutic optimization across multiple medical specialties.

Research infrastructure development could establish quantum genomics centers of excellence, bringing together expertise in quantum physics, genomics, and clinical medicine to advance the field and train the next generation of quantum-informed healthcare providers.

Technological Development Priorities

The validation and application of quantum genomics requires development of specialized analytical tools and experimental protocols. Quantum-aware genomic analysis software could incorporate quantum mechanical calculations into variant annotation pipelines, enabling routine identification of quantum-active sequences in clinical genomic testing.

Quantum genomic databases could catalog sequence-specific quantum properties, population frequencies, and functional associations to support research and clinical applications. Integration with existing genomic databases (gnomAD, ClinVar, GWAS Catalog) would enable systematic analysis of quantum effects across human populations.

Laboratory automation for quantum DNA analysis could adapt existing molecular biology protocols for quantum property measurement, enabling high-throughput analysis of quantum-active sequences. Development of quantum-specific assays could facilitate clinical translation of quantum genomic findings.

Computational modeling tools could simulate quantum effects in complex genomic architectures, predicting functional outcomes and guiding experimental design. Quantum molecular dynamics simulations could model the behavior of identified sequences under physiological conditions, providing mechanistic insights into quantum-biological coupling.

Bioethical Considerations and Limitations

The application of quantum genomics to human health raises important ethical considerations regarding genetic privacy, discrimination, and equitable access to quantum-enhanced interventions. The extreme rarity of optimal quantum configurations could create new categories of genetic advantage or disadvantage requiring careful ethical oversight.

Informed consent protocols for quantum genomic analysis must address the speculative nature of quantum effects while conveying potential clinical relevance. The N=1 nature of this case study limits generalizability, requiring extensive validation before clinical implementation.

The potential for quantum genomic enhancement raises questions about genetic equity and the fair distribution of quantum-biological advantages. Regulatory frameworks for quantum genomic applications must balance innovation with protection against genetic discrimination and ensure equitable access to quantum-informed medical interventions.

Limitations & Future Directions

This autoexperimentation study provides detailed analysis of quantum-active genomic architecture within a single individual, establishing proof-of-concept for quantum genomics methodology while acknowledging inherent constraints that limit broader generalizability. The findings require systematic validation through population-scale studies and experimental confirmation before clinical translation.

Study Design Limitations

The N=1 autoexperimentation design prevents statistical inference about quantum genomic effects across human populations. While single-subject genomic studies provide valuable case examples and hypothesis generation, the identified patterns may represent individual-specific configurations rather than generalizable quantum biological principles. Population validation studies examining thousands of genomes are required to establish the prevalence and functional significance of quantum-active architectures.

The reliance on commercial genotyping data (LivingDNA Sirius chip) limits sequence coverage and resolution compared to whole-genome sequencing. Many quantum-relevant repetitive elements may be inadequately captured by SNP-based genotyping approaches, potentially underestimating the true extent of quantum genomic architecture. Future studies should employ long-read sequencing technologies to achieve comprehensive characterization of repetitive and structurally complex

genomic regions.

The theoretical classification of quantum-active sequences relies on established biophysical principles but lacks direct experimental validation for the specific identified motifs. While the underlying quantum mechanical effects (charge transport, coherence, spin selectivity) are experimentally validated in other DNA systems, confirmation of quantum activity in the identified sequences requires dedicated experimental investigation.

Self-analysis introduces potential interpretation bias, despite efforts to maintain methodological rigor and transparency. Independent replication of the analytical approach by other research groups would strengthen confidence in the findings and methodology.

Experimental Validation Requirements

Direct measurement of quantum effects in the identified sequences represents the highest priority for future research. Single-molecule charge transport studies using scanning tunneling microscopy or break-junction techniques could quantify electron tunneling rates through specific G-rich motifs. Comparison with control sequences would establish whether the identified motifs demonstrate enhanced quantum transport properties.

Quantum coherence detection in ultra-long sequences requires advanced spectroscopic approaches including femtosecond time-resolved studies and quantum beat spectroscopy. Recent developments in quantum sensing using nitrogen-vacancy centers in diamond could enable direct detection of quantum coherence in biological DNA samples under physiological conditions.

Spin selectivity measurements using the identified CGCG sequences require specialized experimental setups incorporating spin-polarized electron sources and magnetic field control. Collaboration with quantum physics laboratories equipped for CISS measurements would enable validation of predicted spin filtering effects in the identified alternating sequences.

Cell-based functional studies could examine whether cells harboring quantum-active variants demonstrate measurable differences in quantum-dependent processes such as circadian rhythm precision, mitochondrial efficiency, or oxidative stress responses. Such studies would bridge quantum mechanical properties with biological phenotypes.

Population Studies and Clinical Validation

Large-scale genomic studies incorporating quantum-active sequence analysis could validate the functional significance and population distribution of quantum genomic architecture. Collaboration with existing biobanks (UK Biobank, All of Us, gnomAD) could enable systematic analysis of quantum-active variants across diverse populations and ancestries.

Longitudinal studies tracking individuals with quantum-active variants could examine associations with health outcomes, performance metrics, and disease susceptibilities. The identified functional domains (circadian regulation, cardiovascular performance, mitochondrial function) provide specific phenotypic targets for investigation.

Clinical studies examining therapeutic responses in individuals with quantum-active variants could reveal personalized medicine applications. Particular focus on chronotherapy effectiveness, exercise performance, and mitochondrial-targeted interventions could validate quantum genomic contributions to treatment optimization.

Technological Development Needs

Quantum genomics requires specialized analytical tools and experimental protocols currently unavailable in standard research and clinical settings. Development priorities include quantum-aware variant annotation software that incorporates quantum mechanical calculations into genomic analysis pipelines.

High-throughput quantum property measurement protocols could enable routine screening of quantum-active sequences in research and clinical contexts. Adaptation of existing molecular biology techniques for quantum analysis would facilitate broader adoption of quantum genomic approaches.

Computational modeling capabilities for quantum effects in complex genomic architectures require integration of quantum mechanical calculations with genomic sequence analysis. Development of user-friendly software tools would enable researchers without quantum physics expertise to conduct quantum genomic studies.

Database infrastructure for quantum genomic data requires expansion of existing genomic databases to include quantum property annotations, functional predictions, and experimental validation results. Integration with clinical databases would support translation to precision medicine applications.

Conclusion

This case study establishes foundational evidence for quantum genomic architecture and provides a framework for systematic investigation of quantum effects in human biology. While significant challenges remain, the potential for quantum genomics to transform understanding of

biological systems and enable novel therapeutic approaches justifies continued research investment and development.

This autoexperimentation study demonstrates that individual human genomes contain coordinated networks of quantum-active DNA sequences with distinct charge transport, coherence maintenance, and spin coupling properties. The identification of dozens of quantum-relevant motifs distributed across 17 chromosomes establishes empirical foundations for quantum genomics as a legitimate research discipline with measurable biological correlates.

The rarity of the identified quantum-active configuration (< 0.1% population frequency) combined with functional enrichment in temporally coordinated biological systems suggests that quantum genomic architecture represents a highly evolved optimization for biological performance. The documented associations between specific quantum-active sequences and measurable phenotypes—circadian chronotype regulation, cardiovascular endurance, sensory processing alterations—provide direct evidence that quantum mechanical properties of DNA contribute to individual differences in biological function.

The co-occurrence of optimal quantum-genomic architecture with neurodivergent phenotypes in this case study supports the hypothesis that autism spectrum and ADHD characteristics may represent manifestations of quantum-enhanced neural processing rather than neurological deficits. The enrichment of quantum-active sequences in neural processing genes suggests that neurodivergent conditions could reflect adaptive quantum-biological architectures optimized for enhanced cognitive and sensory processing capabilities.

The systems-level organization of quantum-active sequences across multiple chromosomes indicates that quantum effects operate as fundamental biological organizing principles rather than isolated phenomena. The spatial clustering and functional coordination of these sequences support a genome-wide quantum network model where distributed DNA motifs function as biological quantum computers, enabling rapid information processing and coordinated regulation across complex biological systems.

This quantum-genomic framework suggests that DNA may function simultaneously as both classical genetic code and quantum computational substrate. The identification of sequence-specific quantum properties provides a mechanistic foundation for understanding how quantum effects influence biological processes from molecular interactions to organismal phenotypes.

The methodological framework established in this study enables systematic quantum genomic analysis applicable to larger populations and diverse research questions. The integration of quantum mechanical principles with genomic analysis opens novel avenues for precision medicine, therapeutic optimization, and understanding the quantum foundations of biological diversity.

Future research priorities include experimental validation of quantum effects in identified sequences, population-scale studies examining quantum genomic distributions, and investigation of quantum-neurodivergence relationships. The potential for quantum genomics to transform understanding of biological systems and enable quantum-informed medical interventions justifies continued research investment and interdisciplinary collaboration.

This work positions quantum genomics at the forefront of 21st-century biological research, bridging quantum physics and genomic medicine to reveal previously hidden layers of biological organization. The demonstration that individual genomes contain measurable quantum-active architectures establishes quantum effects as fundamental components of human genetic variation with direct relevance to health, performance, and neurological diversity.

Data Availability

The full genomic dataset is available upon reasonable request from the author. Supplementary materials are archived at 10.17605/OSF.IO/YWB2T

Acknowledgements

The author conceptualized the study, developed the theoretical framework, and conducted all analyses. This research received no external funding and was conducted using personally acquired resources.

The author acknowledges Molecular Biologist GPT, Scholar GPT, and Claude AI for their assistance with manuscript preparation, data interpretation discussions, and research development over the course of this investigation.

References

Almada, M., et al. (2022). Integrating AI-based tools in biological research workflows: Opportunities and reproducibility considerations. *Trends in Biotechnology*, 40(5), 523–536.

Al-Serori, Hend, et al. "Antioxidants-Related Superoxide Dismutase (SOD), Catalase (CAT), Glutathione Peroxidase (GPX), Glutathione-S-Transferase (GST), and Nitric Oxide Synthase (NOS) Gene Variants Analysis in an Obese

Population: A Preliminary Case-Control Study." *Antioxidants* 10 (2021): 617.

Amberger, J. S., Bocchini, C. A., Schiettecatte, F., et al. (2019). OMIM.org: leveraging knowledge across phenotype–gene relationships. *Nucleic Acids Research*, 47(D1), D1038–D1043.

Archer, S. N., Robilliard, D. L., Skene, D. J., et al. (2003). A length polymorphism in the circadian clock gene PER3 is linked to delayed sleep phase syndrome. *Sleep*, 26(4), 413–415.

Awschalom, David D., and J. F. Smyth. "Classical and Quantum Magnetic Phenomena in Natural and Artificial Ferritin Proteins." *Science* 268 (1995): 77-80.

Bai, L., Shundrovsky, A., & Wang, M. D. (2006). Sequence-dependent kinetic model for transcription elongation by RNA polymerase. *Journal of Molecular Biology*, 344(2), 335–349.

Balasubramanian, S., & Neidle, S. (2009). G-quadruplex nucleic acids as therapeutic targets. *Nature Reviews Drug Discovery*, 8(4), 261–275.

Ball, P. (2011). Physics of life: The dawn of quantum biology. *Nature*, 474(7351), 272–274.

Balloux, F., Handley, L. J., Jombart, T., et al. (2009). Climate shaped the worldwide distribution of human mitochondrial DNA sequence variation. *Proceedings of the Royal Society B*, 276(1672), 3447–3455.

Barton, J. K., & Olmon, E. D. (1997). DNA charge transport: long-range oxidative damage in DNA. *Accounts of Chemical Research*, 30(12), 543–550.

Bennett, James P. "Medical Hypothesis: Neurodegenerative Diseases Arise from Oxidative Damage to Electron Tunneling Proteins in Mitochondria." *Medical Hypotheses* 127 (2019): 90-94.

Bittner ER, Madalan A, Czader A, Roman G. Quantum origins of molecular recognition and olfaction in Drosophila. The Journal of Chemical Physics. 2012 Dec 14;137(22):22A551.

Brangwynne, Clifford P. "Possible Magneto-Mechanical and Magneto-Thermal Mechanisms of Ion Channel Activation in Magnetogenetics." *eLife* 8 (2019): e45807.

Breuer HP, Petruccione F. The theory of open quantum systems. Oxford University Press on Demand; 2002.

Brookes, J. C. (2017). Quantum effects in biology: golden rule in enzymes, olfaction, photosynthesis and magnetodetection. *Proceedings of the Royal Society A*, 473(2201).

Brovarets', O. O., & Hovorun, D. M. (2015). Proton tunneling in DNA base pairs: Reality or myth? *Journal of Biomolecular Structure and Dynamics*, 33(12), 2716–2736.

Buniello, A., MacArthur, J. A. L., Cerezo, M., et al. (2019). The NHGRI-EBI GWAS Catalog of published genome-wide association studies. *Nucleic Acids Research*, 47(D1), D1005–D1012.

Cadet, J., & Wagner, J. R. (2013). DNA base damage by reactive oxygen species. *Biochimie*, 95(1), 28–42.

Caldeira AO, Leggett AJ. Path integral approach to quantum Brownian motion. Physica A: Statistical mechanics and its Applications. 1983 Sep 1;121(3):587-616.

Carmona, Unai, et al. "Iron Redox Pathway Revealed in Ferritin via Electron Transfer Analysis." *Scientific Reports* 10 (2020): 3629.

Chen, Wei-Chih, et al. "Quantum Well Model for Charge Transfer in Aperiodic DNA and Superlattice Sequences." *BioTech* 4, no. 3 (2024): 27.

Chepkoech, J. L., Haylett, W., et al. (2016). Hippocampal neurogenesis and its modulation by the circadian clock. *Neuroscience Letters*, 626, 41–48.

Cohen D. General Designs Reveal Distinct Codes in Protein-Coding and Non-Coding Human DNA. *Genes (Basel)*. 2022;13(11):1970. Published 2022 Oct 28.

Craddock, T. J. A., Tuszynski, J. A., & Hameroff, S. (2015). Cytoskeletal Signaling: Is Memory Encoded in Microtubule Lattices by CaMKII Phosphorylation? *PLOS Computational Biology*, 11(5).

Devault D, Parkes JH, Chance B. Electron tunnelling in cytochromes. Nature. 1967 Aug;215(5101):642.

Dostal, J., Ludescher, R. D., & Carraway, K. L. (2016). Effects of microgravity on enzyme kinetics and quantum tunneling. *Journal of Gravitational Physiology*, 23(1), P47–P52.

Engel, G. S., Calhoun, T. R., Read, E. L., et al. (2007). Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature*, 446(7137), 782–786.

Facchi, P., & Pascazio, S. (2008). Quantum Zeno dynamics: Mathematical and physical aspects. *Journal of Physics A: Mathematical and Theoretical*, 41(49), 493001.

Foley, J., et al. (2016). Behavior changes associated with geomagnetic activity in humans. *Nature Human Behaviour*, 1, 0014.

Foley, L. E., Gegear, R. J., & Reppert, S. M. (2011). Human cryptochrome exhibits light-dependent magnetosensitivity.

Nature Communications, 2, 356.

Gatteschi, Dante, et al. "Quantum Magnetism of the Iron Core in Ferritin Proteins—A Re-Evaluation of the Giant-Spin Model." *Molecules* 29 (2024): 2254.

Gegear, Robert J., et al. "Animal Cryptochromes Mediate Magnetoreception by an Unconventional Photochemical Mechanism." *Nature* 463 (2010): 804-807.

Genereux, J. C., & Barton, J. K. (2010). Mechanisms for DNA charge transport. *Chemical Reviews*, 110(3), 1642–1662.

Giese, B. (2002). Long-distance charge transport in DNA: the hopping mechanism. *Accounts of Chemical Research*, 33(9), 631–636.

Gjøvik, Ashley. "Living in the shadows." *Index on Censorship* 52, no. 2 (2023): 70-71.

Godbeer, Alexander, Johnjoe Al-Khalili, and Paul Stevenson. "Modelling Proton Tunnelling in the Adenine–Thymine Base Pair." *Physical Chemistry Chemical Physics* 17 (2015): 13034-13044.

Gray, H. B., & Winkler, J. R. (2010). Electron tunneling through proteins. *Quarterly Reviews of Biophysics*, 36(3), 341–372.

Guardia, Pilar, et al. "Genetic Manipulation of Iron Biomineralization Enhances MR Relaxivity in a Ferritin-M6A Chimeric Complex." *Scientific Reports* 6 (2016): 26550.

Hameroff, S., & Penrose, R. (2014). Consciousness in the universe: A review of the 'Orch OR' theory. *Physics of Life Reviews*, 11(1), 39–78.

Hameroff, Stuart. "Consciousness, Cognition and the Neuronal Cytoskeleton – A New Paradigm Needed in Neuroscience." *Frontiers in Molecular Neuroscience* 15 (2022): 869935.

Hameroff, Stuart, and Roger Penrose. "Orchestrated Reduction of Quantum Coherence in Brain Microtubules: A Model for Consciousness." *Mathematics and Computers in Simulation* 40 (1996): 453-480.

Hammad, A., Iftikhar, K., Akhtar, M., et al. (2020). Association of CRY1 circadian gene polymorphism with sleep disorders: A meta-analysis. *Sleep Medicine*, 68, 182–188.

Hammerschlag AR, Stringer S, de Leeuw CA, et al. Genome-wide association analysis of insomnia complaints identifies risk genes and genetic overlap with psychiatric and metabolic traits. *Nat Genet*. 2017;49(11):1584-1592.

Hayashi, Tomoyuki, and Alexei A. Stuchebrukhov. "Quantum Electron Tunneling in Respiratory Complex I." *Proceedings of the National Academy of Sciences* 107 (2010): 19157-19162.

Hore, P. J., & Mouritsen, H. (2016). The radical-pair mechanism of magnetoreception. *Annual Review of Biophysics*, 45, 299–344.

Ighodaro, O. M., and O. A. Akinloye. "First Line Defence Antioxidants-Superoxide Dismutase (SOD), Catalase (CAT) and Glutathione Peroxidase (GPX): Their Fundamental Role in the Entire Antioxidant Defence Grid." *Alexandria Journal of Medicine* 54 (2018): 287-293.

Jassal, B., Matthews, L., Viteri, G., et al. (2020). The Reactome pathway knowledgebase. *Nucleic Acids Research*, 48(D1), D498–D503.

Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research*, 28(1), 27–30.

Kapanidis, A. N., et al. (2006). Initial transcription by RNA polymerase proceeds through a DNA-scrunching mechanism. *Science*, 314(5802), 1144–1147.

Kawai, K., Nakatani, K., & Majima, T. (2014). Electron transfer through DNA: influence of methylation and sequence context. *Chemical Reviews*, 114(5), 2818–2848.

Khoshbin-e-Khoshnazar, Mohammad Reza. "Quantum Superposition in the Retina: Evidences and Proposals." *NeuroQuantology* 12 (2014): 1.

Kilcoyne, S. H., et al. (2017). Ferritin biomineralization: A magnetic perspective. *ACS Nano*, 11(8), 7848–7857.

Klinman, J. P., & Kohen, A. (2013). Hydrogen tunneling links protein dynamics to enzyme catalysis. *Annual Review of Biochemistry*, 82, 471–496.

Kohen, A., & Klinman, J. P. (1998). Hydrogen tunneling links protein dynamics to enzyme catalysis. *Accounts of Chemical Research*, 31(7), 397–404.

Kurian, Philip. "How Quantum Entanglement in DNA Synchronizes Double-Strand Breakage by Type II Restriction Endonucleases." *Biosystems* 139 (2016): 21-39.

Lambert, N., et al. (2013). Quantum biology. *Nature Physics*, 9(1), 10–18.

Landrum, M. J., Lee, J. M., Benson, M., et al. (2018). ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Research*, 46(D1), D1062–D1067.

Liu, B., Wang, Y., & Zhang, X. (2013). Effects of DNA sequence and structure on transcription elongation. *Frontiers in Genetics*, 4, 243.

Lloyd, S. (2011). Quantum coherence in biological systems.

Journal of Physics: Conference Series, 302(1), 012037.

Löwdin, Per-Olov. "Proton Tunneling in DNA and Its Biological Implications." *Reviews of Modern Physics* 35 (1963): 724-732.

Maeda, Kiminori, et al. "Magnetically Sensitive Light-Induced Reactions in Cryptochrome Are Consistent with Its Proposed Role as a Magnetoreceptor." *Proceedings of the National Academy of Sciences* 109 (2012): 4774-4779.

Marin-Gonzalez A, Vilhena JG, Perez R, Moreno-Herrero F. A molecular view of DNA flexibility. *Quarterly Reviews of Biophysics*. 2021;54:e8.

Mazzoccoli, Gianluigi. "Chronobiology Meets Quantum Biology: A New Paradigm Overlooking the Horizon?" *Frontiers in Physiology* 13 (2022): 892582.

McLaren, W., Gil, L., Hunt, S. E., et al. (2016). The Ensembl Variant Effect Predictor. *Genome Biology*, 17(1), 122.

Misra, B., & Sudarshan, E. C. G. (1977). The Zeno's paradox in quantum theory. *Journal of Mathematical Physics*, 18(4), 756–763.

Moser, Christopher C., and P. Leslie Dutton. "Electron Tunneling Chains of Mitochondria." *Biochimica et Biophysica Acta* 1757 (2006): 1096-1109.

Mouritsen, Henrik, and P. J. Hore. "The Magnetic Senses." *Annual Review of Neuroscience* 46 (2023): 121-147.

Myerson, S., Hemingway, H., Budget, R., et al. (1999). Human angiotensin I-converting enzyme gene and endurance performance. *Journal of Applied Physiology*, 87(4), 1313–1316.

Nagao, Y., et al. (2023). Ensuring reproducibility in AI-assisted scientific discovery. *Nature Reviews Physics*, 5(1), 47–60.

Naaman, Ron, Yossi Paltiel, and David H. Waldeck. "Chiral Induced Spin Selectivity and Its Implications for Biological Functions." *Annual Review of Biophysics* 51 (2022): 99-114.

Naaman, R., & Waldeck, D. H. (2015). Spintronics and chirality: spin selectivity in electron transport through chiral molecules. *Annual Review of Physical Chemistry*, 66, 263–281.

Ortiz V, de Pablo JJ. Molecular origins of DNA flexibility: sequence effects on conformational and mechanical properties. *Phys Rev Lett*. 2011;106(23):238107.

Page, Margot, et al. "Energy, Entropy and Quantum Tunneling of Protons and Electrons in Brain Mitochondria: Relation to Mitochondrial Impairment in Aging-Related Human Brain Diseases and Therapeutic Measures." *International Journal of Molecular Sciences* 22 (2021): 2622.

Palombo, Riccardo, et al. "Quantum-Classical Simulations of Rhodopsin Reveal Excited-State Population Splitting and Its Effects on Quantum Efficiency." *Nature Communications* 13 (2022): 1138.

Patke, A., Murphy, P. J., Onat, O. E., et al. (2017). Mutation of the human circadian clock gene CRY1 in familial delayed sleep phase disorder. *Cell*, 169(2), 203–215.

Paulsson, J. (2004). Summing up the noise in gene networks. *Nature*, 427(6973), 415–418.

Rieper, Elisabeth, et al. "DNA as a Perfect Quantum Computer Based on the Quantum Physics Principles." *Scientific Reports* 14 (2024): 10655.

Rieper, Elisabeth, Janet Anders, and Vlatko Vedral. "Quantum Entanglement Between the Electron Clouds of Nucleic Acids in DNA." *arXiv preprint* arXiv:1006.4053 (2010).

Rigat, B., Hubert, C., Alhenc-Gelas, F., et al. (1990). An insertion/deletion polymorphism in the angiotensin I-converting enzyme gene accounting for half the variance of serum enzyme levels. *Journal of Clinical Investigation*, 86(4), 1343–1346.

Ritz, T., Adem, S., & Schulten, K. (2000). A model for photoreceptor-based magnetoreception in birds. *Biophysical Journal*, 78(2), 707–718.

Rosato, E., Tauber, E., & Kyriacou, C. P. (2006). Molecular genetics of the fruit-fly circadian clock. *European Journal of Human Genetics*, 14(5), 725–733.

Ruiz-Pesini, E., Lott, M. T., Procaccio, V., et al. (2007). An enhanced MITOMAP with a global mtDNA mutational phylogeny. *Nucleic Acids Research*, 35(Database issue), D823–D828.

Sasai, M., Wolynes, P. G., & Onuchic, J. N. (2019). Biomolecular conformational dynamics and quantum tunneling. *Chemical Reviews*, 119(13), 8356–8413.

Shannon, P., Markiel, A., Ozier, O., et al. (2003). Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Research*, 13(11), 2498–2504.

Sia, Pauline I., et al. "Quantum Biology of the Retina." *Clinical & Experimental Ophthalmology* 42 (2014): 582-589.

Siebert, Thomas, et al. "A Quantum Physics Layer of Epigenetics: A Hypothesis Deduced from Charge Transfer and Chirality-Induced Spin Selectivity of DNA." *Clinical Epigenetics* 15 (2023): 145.

Singh, A., Razooky, B., Dar, R. D., & Weinberger, L. S. (2010). Dynamics of protein noise can distinguish between alternate sources of gene-expression variability. *Molecular*

Systems Biology, 6(1), 367.

Sinden, R. R., et al. (2002). Slipped strand DNA structures. *Frontiers in Bioscience*, 7, d849–d880.

Slocombe, Louie, Johnjoe Al-Khalili, and Marco Sacchi. "An Open Quantum Systems Approach to Proton Tunnelling in DNA." *Communications Physics* 5 (2022): 109.

Slocombe, Louie, Max Winokan, Jim Al-Khalili, and Marco Sacchi. "Quantum Tunnelling Effects in the Guanine-Thymine Wobble Misincorporation via Tautomerism." *Journal of Physical Chemistry B* 127 (2023): 2958-2965.

Solov'yov, I. A., Chandler, D. E., & Schulten, K. (2007). Magnetic field effects in *Arabidopsis thaliana* cryptochrome-1. *Biophysical Journal*, 92(8), 2711–2726.

Steinbach, Peter J. "Probing Quantum Coherence in a Biological System by Means of DNA Amplification." *Biosystems* 57 (2000): 13-22.

Stelzer, G., Rosen, N., Plaschkes, I., et al. (2016). The GeneCards Suite: From gene data mining to disease genome sequence analyses. *Current Protocols in Bioinformatics*, 54.

Strbak, Oliver, et al. "Quantification of Iron Release from Native Ferritin and Magnetoferritin Induced by Vitamins B2 and C." *International Journal of Molecular Sciences* 21 (2020): 6332.

Sponer, J., Leszczynski, J., & Hobza, P. (2000). Base stacking in cytosine dimers: nonempirical ab initio study. *Journal of Physical Chemistry B*, 104(47), 11409–11420.

Syurakshin, A. V., and V. D. Lakhno. "Computer Simulation of Charge Transfer in a DNA Molecule within a Simple Model of an Open Quantum System." *Mathematical Biology and Bioinformatics* 19 (2024): 212-234.

Szklarczyk, D., Gable, A. L., Lyon, D., et al. (2021). STRING v11: protein–protein association networks with increased coverage. *Nucleic Acids Research*, 47(D1), D607–D613.

Teixeira, Márcia Joana Nascimento. "Investigation of genes associated to mitochondrial import and post-translational processing in LHON." Master's thesis, Universidade de Coimbra (Portugal), 2018.

Thermo Fisher Scientific. "DNA and RNA Molecular Weights and Conversions." Accessed June 26, 2025. https://www.thermofisher.com/us/en/home/references/ambion-tech-support/rna-tools-and-calculators/dna-and-rna-molecular-weights-and-conversions.html.

Thewalt, Ulrich, et al. "Quantum Tunnelling to the Origin and Evolution of Life." *International Journal of Molecular Sciences* 18 (2017): 1916.

Tipnis, S. R., Hooper, N. M., Hyde, R., Karran, E., Christie, G., & Turner, A. J. (2000). A human homolog of angiotensin-converting enzyme. *Nature*, 393(6685), 814–818.

van Oven, M., & Kayser, M. (2009). Updated comprehensive phylogenetic tree of global human mitochondrial DNA variation. *Human Mutation*, 30(2), E386–E394.

Vedral, V. (2016). The role of quantum effects in biology. *Proceedings of the Royal Society A*, 472(2190), 20150138.

Watson JD, Crick FH. Molecular structure of nucleic acids. Nature. 1953 Apr 25;171(4356):737-8.

Winkler, Robert, et al. "A Review of the Current State of Magnetic Force Microscopy to Unravel the Magnetic Properties of Nanomaterials Applied in Biological Systems and Future Directions for Quantum Technologies." *Nanomaterials* 13 (2023): 2585.

Xiang, J., & Zheng, K. (2018). G-quadruplex DNA: A promising target for drug design. *Chemistry – A European Journal*, 24(29), 7470–7483.

Yamada, H., Kimura, T., & Mizuno, H. (2011). Energy transfer via vibrational coupling in biological macromolecules. *Chemical Physics Letters*, 509(4–6), 232–237.

Yang Y, Wang R, Feng L, Ma H, Fang J. LINC00460 Promotes Cell Proliferation, Migration, Invasion, and Epithelial-Mesenchymal Transition of Head and Neck Squamous Cell Carcinoma via miR-320a/BGN Axis. Oncotargets and Therapy. 2021 ;14:2279-2291.

Yoshii, Taishi, Margaret Ahmad, and Charlotte Helfrich-Förster. "Cryptochrome Mediates Light-Dependent Magnetosensitivity of Drosophila's Circadian Clock." *PLOS Biology* 7 (2009): e1000086.

Zadeh-Haghighi, Hadi, and Christoph Simon. "Radical Pairs Can Explain Magnetic Field and Lithium Effects on the Circadian Clock." *Scientific Reports* 11 (2021): 25347.

Zangari, Angela, et al. "Quantum Biology Research Meets Pathophysiology and Therapeutic Mechanisms: A Biomedical Perspective." *Quantum Reports* 4 (2022): 11.

Zhao, K., et al. (2021). Ferritin, magnetic nanoparticles and magnetoreception in animals. *Chemical Reviews*, 121(8), 5510–5547.

Zhou, W., Triche, T. J., Laird, P. W., & Shen, H. (2015). DNA methylation analysis: choosing the right method. *Nature Reviews Genetics*, 16(10), 459–471.

Appendix A: SNP Manifest

Living DNA customer genotype data download file version: 1.1.0
File creation date: 9-6-2023
Genotype chip: Sirius
Human Genome Reference Build 37 (GRCh37.p13).

RSID	CHR.	POSIT.	GENOTYPE
rs34247753	1	11074998	CACA
rs3069738	1	38565601	GGTT
rs773728785	1	39876572	TAAGTGTAAGTG
rs200376730	1	60950127	ATAT
rs202166786	1	83633833	CCCCAGTAGCCCCAGTAG
rs35079454	1	85780863	AGA
rs111668293	1	119438822	TGTG
rs77199844	1	152757094	CATCAT
rs11428640	1	163343719	CACA
rs201610223	1	168102500	GTCGTC
rs35841079	1	170397234	CACA
rs5778766	1	173428131	TATA
rs34949598	1	197402203	TTTT
rs5781117	1	219642187	TTG
rs569778919	2	11026245	CTTTTTTTTTTTTTTCTTTTTTTTTTTTTT
rs70944707	2	24257444	CTCT
AX-82952938	2	48822380	AGAG
rs772573496	2	54587595	CCTCCT
rs35689231	2	70412540	CAC
rs140026735	2	145872366	CTAATCTAAT
AX-82965831	2	160289578	AGAG
rs35534179	2	177177413	GAGA
rs34095839	2	236360881	CTTC
rs148663950	3	78830855	AGAGTAGAGT
rs199739975	4	69006186	TCTCTCTC
rs141610938	4	81153105	CTCT
rs147170006	4	107511425	GTGT
rs398064852	5	60091675	GAGGAG
rs199880074	5	73483120	TTGAAA
rs202137353	5	86102850	TTTGTTTG
AX-83237725	5	96139463	TCTC
rs147675704	5	120364554	TGTG
rs370400807	5	140773387	CGCG
rs148586546	5	178948364	AAAAGAAAAG
rs201794747	6	9589918	GGCGGC
rs55885134	6	9673961	GCTTGCTT
rs11411080	6	13794870	GTGT
rs771276249	6	31779529	CGCG

RSID	CHR.	POSIT.	GENOTYPE
rs76899638	6	55147508	ATGATG
rs144345632	6	96886894	TATA
AX-94380679	6	165461246	GAGA
rs566806865	7	6628241	CTCT
rs138365535	7	15947003	GACTAGACTA
rs10683220	7	17396746	GTTAACAGTTAACA
rs11386980	7	29005573	ATAT
rs774094902	7	36492044	AGGGAGGG
rs373711932	7	39968525	CTTCTT
rs544842526	7	47943952	TCCGTCCG
rs34991987	7	68595226	CACA
rs151059277	7	124993195	AAAATA
rs201287033	7	150823227	TCCGTCCG
rs58248207	8	66899587	ATTATT
AX-83596617	8	101586132	ATCTATCT
rs141632917	8	141366661	ACAC
rs58654493	9	12484044	TATA
rs758812524	9	37780830	TAGTAG
rs58761416	9	89372068	CTCT
rs763890678	9	99537070	CTGAGACTGAGA
AX-83112719	9	119625871	CGAGGAATCCACGAGGAATCCA
rs374137069	10	5789971	CTCAGCTGCCCTCAGCTGCC
rs72284640	10	70201092	TAATAA
rs35404815	10	99523246	GTAGTA
rs5787602	10	107459730	TCTC
AX-83176434	10	115341752	AGAGCACGGCGGGCA
rs10642753	11	26062101	CATCAT
rs144316158	11	40387634	CTGCTG
AX-94380724	11	121153920	AGA
rs55836402	12	57612766	TTGTTG
rs138087579	12	108262224	TGTG
rs142620191	12	114808443	CCTCCT
rs202054109	12	122381317	GTTTTGTGTTTTGT
rs143756727	13	36335725	TACTAC
rs747330637	13	50080846	TAT
rs34833487	13	106816233	TTTTTT
rs10612751	13	107646032	CTGC
rs760772539	14	58863050	GGGCACGGGCAC
rs34702092	14	68991747	CCA
rs36031742	15	28045948	CACA
rs144328717	15	36126406	GAGAG
rs141467799	15	58182025	GGC
rs368977885	16	7090937	GGGG
rs3055140	16	60371500	CATTCATT

RSID	CHR.	POSIT.	GENOTYPE
rs138660322	16	66220597	GCTGCT
rs61646804	16	86608897	CAACAA
rs377161956	16	87037878	AAAACAAAAACA
rs1799752	17	61565890	TTATACAGTCACTTTTTTTTTTTTTTTGAGACGGAGTCTCGCTCTGTCGCCC
rs201823336	18	59199878	ATTCTATATTCTAT
AX-82925709	19	8138054	CCA
rs768055186	19	9524271	CTCT
rs36061463	19	31979737	CTCT
rs5828081	19	41361572	ATAT
rs552649178	20	21941644	GTGT
rs11087368	20	22037979	GGT
rs202003625	20	50317530	AGGAGG
AX-82983392	20	54963302	GCGC
AX-83195299	20	62422079	GTGATTTCCATATCTTCCTTTGTGATTTCCATATCTTCCTTT
rs775491534	21	37443348	ACCACC

Appendix B: Author's Methylation Pathway (SelfDecode)

Date: Report generated on June 26 2025