Deep Time Geology in Santa Clara County

Evidence for Two Billion Years of Active Continental Margin Processes

Authors: Ashley M. Gjøvik, B.S., J.D. (ashleymgjovik@protonmail.com) | **Date**: July 15 2025

Abstract:

Santa Clara County preserves geological evidence spanning two billion years of Earth history, challenging conventional interpretations that describe this region as comprising scattered Mesozoic oceanic fragments. Systematic analysis of isotopic, geochemical, structural, and ecological data reveals active preservation of Precambrian crustal materials and processes rather than passive remnants of Jurassic-Cretaceous oceanic crust.

Recent isotopic studies provide direct evidence for this ancient heritage. Coast Range Ophiolite peridotites yield model ages of ~1.5 Ga. Point Sal peridotites show >20% melt depletion with 187Os/188Os ratios indicating significant ancient signatures preserved in refractory mantle domains.

Modern geochemical conditions match optimal parameters for serpentinization indicating ongoing deep crustal activity rather than passive remnants. Historical ecological patterns, systematic earthquake damage distributions, and organized geochemical signatures reflect surface expression of ancient mantle processes that remain structurally active through fault-controlled circulation systems.

Mercury mineralization, alkali meadow distribution, and integrated hydrothermal circulation demonstrate that billion-year-old serpentinization systems continue operating beneath the urbanized landscape. Analysis reveals systematic transport of blueschist-facies minerals and Franciscan greenstone, with blue amphibole concentrations increasing with depth, indicating ongoing erosion of high-pressure metamorphic systems.

Santa Clara County represents a natural laboratory where deep-time Earth processes span multiple supercontinent cycles and remain directly observable through their continued influence on modern surface environments. The integration of ancient materials with active fault systems creates opportunities to study billion-year geological evolution.

Keywords:

deep time geology, Precambrian, supercontinent cycles, serpentinization, continental margin evolution, Santa Clara County, Silicon Valley, California, Coast Range Ophiolite, Franciscan Complex, blueschist metamorphism, hydrothermal systems, isotopic analysis, fault-controlled circulation, active geological systems, Nuna supercontinent, Rodinia, geological mapping

Citation:

Gjøvik, Ashley M. “Deep Time Geology in Santa Clara County: Evidence for Two Billion Years of Active Continental Margin Processes.” *The Journal of Decolonized Ecology and Evolution* 1, no. 1 (2025). DOI 10.5281/zenodo.15892989

Table of Contents

Introduction

The San Francisco Bay Area represents one of the most geologically complex terrains in North America, shaped by crustal interaction, oceanic accretion, mantle exposure, and plate reorganization. Its current configuration results from repeated accretionary and obductionary events that assembled ancient oceanic and continental fragments, culminating in modern exposure of deep-sea lithologies on continental surfaces.

Santa Clara County occupies a critical position by preserving geological relationships that span from Archean mantle processes through multiple supercontinent cycles to modern transform tectonics. The region preserves evidence of complete Wilson cycles, with mineral assemblages recording distinct phases of Earth's deep-time evolution.

The evidence supporting this interpretation includes: isotopic signatures documenting billion-year model ages with direct connections to Nuna/Columbia supercontinent processes; systematic surface patterns indicating active subsurface processes rather than passive geological remnants; integrated circulation systems maintaining continuous access to ancient materials through fault-controlled networks; and ongoing chemical activity demonstrating that ancient systems remain structurally active and continue to influence modern surface environments.

This analysis employs systematic integration of isotopic, geochemical, ecological, and structural evidence to develop a comprehensive framework explaining Santa Clara County's acknowledged geological complexities. The convergence of independent analytical approaches provides robust support for the deep-time interpretation while demonstrating the limitations of conventional models that treat the region as a collection of disconnected geological fragments.

The study demonstrates that conventional geological mapping, when combined with deep-time analysis, reveals Santa Clara County as a natural laboratory preserving direct evidence of Earth's deep-time evolution. The ongoing operation of billion-year-old processes offers unprecedented opportunities to study continental margin development and the preservation of ancient geological systems.

The following sections present detailed evidence supporting this framework, demonstrating how systematic analysis of regional geological patterns reveals Santa Clara County as a unique window into processes spanning Earth's supercontinent cycles and their continued influence on modern surface environments. A separate forthcoming paper will also discuss the deep-time biological implications of this present study.

Santa Clara Valley: The Valley of Heart's Delight

Santa Clara County was built upon what geologists now recognize as some of the most complex and chaotic geological terrain in North America. The central portion of Santa Clara County was fertile agricultural flatland once referred to as "The Valley of Heart's Delight," yet this same area sits atop a geological structure so complex and mysterious that there remains no consensus on the underlying geology, only consensus that there are multiple, chaotic formations and that there is generally no consensus (Langenheim, 2015; Santa Clara Co., 2012).

Since at least 1769, residents and travelers have described the Santa Clara Valley in extraordinary terms. Writer Bayard Taylor, traveling in 1859, described the valley as "unlike anything else in the world; with a beauty so new and dazzling, that all ordinary comparisons are worthless" (1951). Land surveyor Tracy (1853) described a valley "remarkable for its beauty and fertility" (Beller et al., 2010).

The 1881 History of Santa Clara County describes the soil in Santa Clara Valley as producing "a crop of marvelous profusion and excellence" (Munro-Fraser, 1881). The U.S. government reported that orchard crops from Santa Clara Valley "were of unusually high quality" with "beautiful fruit" that created "international acclaim" (USDA, 2015).

What made this agricultural success particularly remarkable was its consistency and exceptional quality across diverse crops. The valley supported everything from grain fields to world-famous orchards, with fruit quality that set international standards. This extraordinary fertility across such a broad range of agricultural products suggests something systematic about the underlying soil chemistry and geological processes.

Serpentinization reactions release Mg^{2+}, Ca^{2+}, and OH^- ions into circulating groundwater, creating naturally alkaline conditions (pH 8-11) optimal for nutrient availability. The process also releases trace elements including boron, molybdenum, and selenium which are essential for plant growth but typically limiting in California soils.

As this mineral-rich groundwater reaches surface through springs and diffuse seepage, it deposits these nutrients across the valley floor through irrigation and seasonal flooding. The resulting soil chemistry (high cation exchange capacity, optimal pH buffering, and trace element supplementation) created the systematic agricultural advantage observed across diverse crop types, explaining both the consistency and exceptional quality that set international standards.

Today, soil across Santa Clara County generally has high electrical conduction (strongly saline), contains alkaline water, has high sodium absorption and high cation exchange, and has poor drainage (Walkinshaw et al., 2023). This alkaline, mineral-rich chemistry that once produced extraordinary crops reflects systematic geochemical processes.

The current valley floor represents a relatively recent development in a much longer geological story. The 10-17 Ma topography of what is now Santa Clara Valley probably resembled the present-day Santa Cruz Mountains or Diablo Range, with broad areas of shallow marine deposition and local areas of deep marine deposition (Langenheim, 2015). The transformation from mountainous terrain to fertile valley occurred through processes operating on deep-time timescales.

Continental margins experience relatively intense geological activity compared to most places over time. California's long, complex, and active geologic history explains the diversity of rock types and the variety of geologic processes that have acted on these rocks, leading to the formation of an unusual diversity of mineral deposits (Dellinger, 1989). Santa Clara County sits at the intersection of these continental margin processes, preserving a record of geological evolution that extends deep into Earth's history.

Deep Time Geology

The Framework: Solving Santa Clara County's Geological Mysteries

Santa Clara County represents one of the most geologically complex and scientifically significant regions in California, yet conventional geological frameworks have repeatedly failed to explain the systematic patterns observed. While there's no consensus on the underlying geology, there is consensus that there are multiple, chaotic formations, complexes, and sediments—including overlaid and intersecting groups of rock referred to as the Franciscan Complex, Coast Range ophiolite, Sierra Azul block, Santa Clara Formation, Temblor Formation, and the Great Valley Sequence (Langenheim, 2015).

Researchers openly acknowledge fundamental gaps in understanding. "The distribution of Santa Clara gravels...is puzzling" and "no one tectonic model completely explains the history of Santa Clara Valley" (Langenheim, 2015). More broadly, "Little is known about the early geologic history of California" (Dellinger, 1989). Earthquake studies reveal systematic damage patterns where "The origin of damage in this zone is elusive" as Santa Clara does not have mapped faults (Brabb and Hanna, 1981; Carter and Jachens, 1993).

When shallow-time models repeatedly fail to explain systematic patterns, the answer is usually that the patterns are older and more systematic than assumed. The complexity observed in Santa Clara County requires time and process scales that conventional Mesozoic timelines simply cannot provide.

This study represents the first systematic attempt to explain Santa Clara County's acknowledged geological mysteries by working backwards from what we observe to determine what processes and timeframes could actually produce these results. The answer: two billion years of preserved crustal evolution.

Isotopic Evidence for Ancient Mantle Heritage

Isotopic analysis acts like a geological fingerprint, revealing when and how rocks formed by measuring the ratios of different atomic versions (isotopes) of elements within minerals. When rocks undergo major geological processes (like melting, crystallization, or alteration) they acquire distinctive isotopic signatures that persist for billions of years. These signatures allow geologists to trace materials back to their original formation environments and dates, even when the rocks have been heavily modified by later events.

For Santa Clara County's geological materials, isotopic evidence provides the most direct proof that we're dealing with far more ancient processes than conventional models recognize. Rather than simple 160-million-year-old oceanic fragments, the isotopic signatures reveal billion-year-old mantle materials that preserve records of Earth's deep-time evolution.

Direct Evidence for Billion-Year Heritage

Recent isotopic studies provide compelling proof that Santa Clara County's geological materials preserve ancient signatures spanning back to supercontinent cycles. Coast Range Ophiolite peridotites yield aluminachron model ages of ~1.5 Ga, with researchers noting this "may correspond to rifting of the Columbia supercontinent" (Choi et al., 2024). This demonstrates that these materials preserve Mesoproterozoic mantle signatures and evidence of processes that occurred during the assembly and breakup of ancient supercontinents.

Model ages differ from crystallization ages. While a rock may have been physically emplaced in its current location relatively recently, model ages reveal when the source material underwent its original differentiation from primitive mantle. A 1.5 billion-year model age means the mantle source experienced major melting events during supercontinent cycles, creating the chemical signatures we observe today.

Multiple Lines of Isotopic Evidence

The Coast Range Ophiolite peridotites are divided into two groups: abyssal and supra-subduction zone types. The abyssal-type peridotites are particularly significant because they represent residues after ≤5% melting of the primitive upper mantle. They are remnants of oceanic lithosphere that were trapped in a subduction setting before being re-melted or modified by later processes.

Critically, these abyssal-type peridotites yield an aluminachron model age of ~1.5 Ga, implying that the Coast Range Ophiolite mantle had experienced episodes of melt extraction before the current ophiolite crust was formed (Choi et al., 2024). This means we're not looking at simple oceanic crust that formed 160 million years ago, but rather ancient mantle materials that preserve evidence of billion-year-old melting events.

Point Sal peridotites provide additional confirmation of ancient heritage, showing >20% melt depletion of a fertile mantle protolith with 187Os/188Os = ~0.129, indicating significant ancient signatures (Snortum & Day, 2020). These peridotites have 187Os/188Os values ranging from 0.1188 to 0.1315. These ratios are crucial for understanding their history.

Osmium isotope ratios (187Os/188Os) act like a geological clock for mantle materials. Very low ratios (like those found here) indicate materials that have been depleted of certain elements for extremely long periods. This is exactly what we'd expect from ancient mantle that underwent major melting events billions of years ago. Further, some refractory abyssal peridotites show very depleted ratios implying long-term preservation of refractory mantle domains.

Evidence of Active, Preserved Systems

The preservation of distinct highly siderophile element patterns, coupled with low 187Os/188Os values, supports a scenario where these peridotites were not substantially rehomogenized by later thermal events, enabling billion-year preservation. In most geological settings, ancient signatures get erased by subsequent heating and chemical exchange. The fact that these ancient signatures survived indicates very specific preservation conditions.

Critically, these studies also document Late Miocene alteration events affecting the peridotites, interpreted as related to tectonic changes and hydration-dehydration reactions in response to changes in Pacific-North American plate motion (Snortum & Day, 2020). This proves these ancient systems remain structurally active and accessible, not passive fossil remnants.

The combination of billion-year model ages with evidence of recent alteration demonstrates that ancient materials continue to interact with modern geological processes. This ongoing activity is exactly what we'd expect if Santa Clara County preserves active deep-time systems rather than isolated fragments.

Implications for Regional Understanding

These aren't 161 Ma oceanic fragments that happened to get emplaced during Mesozoic subduction. They're ancient mantle materials with documented Mesoproterozoic signatures that preserve direct records of supercontinent-scale processes. The isotopic evidence directly contradicts conventional models that treat this region as a collection of relatively young oceanic remnants.

The isotopic signatures provide a direct temporal link between Santa Clara County materials and the deep-time processes that shaped early continental margins during supercontinent cycles.

The preservation of these ancient signatures, combined with evidence for ongoing alteration and access, indicates that Santa Clara County doesn't just contain old rocks. It preserves active systems that have operated continuously since Precambrian time, maintaining their ability to influence modern surface environments through fault-controlled circulation and ongoing serpentinization processes.

Supercontinent Context: Precambrian Continental Margin Processes

Having established that these materials preserve 1.5+ billion year signatures, it becomes necessary to examine the Precambrian geological processes that could produce the systematic patterns observed in Santa Clara County. The temporal framework emerges through analysis of Laurentia's position during key supercontinent cycles and the processes operating along its western continental margin during Mesoproterozoic time.

Nuna/Columbia Assembly (1.8-1.6 Ga): Active Margin Processes

Laurentia first assembled from large fragments of Archean crust around 2.0 to 1.8 Ga during global-scale collisional events, then became the core of the supercontinent Nuna around 1.9-1.6 Ga (Karlstrom et al., 2001). During Nuna assembly, western Laurentia occupied a crucial position in global paleogeography. At 1.6 Ga, E-W-directed collision between the Laurentian and Australian blocks formed a N-S-striking compressional fabric recording Barrovian-type metamorphism (Pourteau et al., 2018).

Evidence confirms that Australia was positioned near North America as part of the Columbia/Nuna supercontinent. Researchers analyzed zircons in Georgetown rocks now in northern Australia and found they matched chemical signatures of similar-aged rocks in northwestern Canada, validating that this chunk of continental crust was transferred from North America to Australia during ancient supercontinent movements (Nordsvan et al., 2018). The west coast of Laurentia (what is now California, Nevada, BC, Yukon) was an active margin, colliding with cratons like Australia, Antarctica, and Siberia.



Alternative reconstructions of the Nuna Supercontinent in 1720 Ma and 1650 Ma. (Gibson and Withnall, 2020).

Rodinia Assembly & Breakup (1.1-0.6 Ga): Margin Development

The Rodinia supercontinent assembled through worldwide orogenic events between 1,300 Ma and 900 Ma, involving all existing continental blocks. Around 1.07 Ga, Laurentia was part of supercontinent Rodinia, rotated approximately 90 degrees clockwise compared with its modern orientation, with East Antarctica and Australia to the north, Siberia to the east, and Baltica and Amazonia to the south (Li et al., 2008).

During Rodinia's breakup (~1.1-1.0 Ga), massive hydrothermal systems activated, and rifting triggered crustal extension, slab rollback, and spreading centers with hydrothermal circulation into old lithosphere. The first major break-up event occurred along the western margin of Laurentia, creating an extensional continental margin from Mexico to Yukon during the breakup of Rodinia about 900–800 Ma, commencing with a series of Precambrian rift basins, followed by the development of a wedge of Paleozoic continental margin sediments.



Reconstruction of Laurentia circa 1.5-1.3 Ga
(Karlstrometal, 2001).

Optimal Conditions for Long-Term Preservation

During the Rodinia breakup (~750–600 Ma), crustal rifting plus slab rollback allowed seawater to percolate into ancient upper mantle, creating serpentinization conditions ideal for preserving ancient materials. This made the rock less dense and chemically reactive, attracting biogenic and silica-rich fluids. These are the perfect conditions for preserving microbial textures, biogenic silica, and opalite veins. Serpentinite is too buoyant to sink deeply into the mantle and tends to stall at the base of subduction zones. Once tectonically sliced into the continental margin, it becomes locked in place.

The preservation mechanism operates through several complementary processes:

- **Density Isolation:** Serpentinized materials (density ~2.6 g/cm³) are significantly less dense than surrounding rocks (~2.8-3.2 g/cm³), preventing subduction recycling and maintaining position in upper crustal levels.
- **Structural Protection:** Location at continental margins provided shelter from intense thermal events that would have destroyed these materials in cratonic interiors.
- **Continuous Hydration**: Ongoing serpentinization reactions consume potential thermal energy that might otherwise drive dehydration and destruction.
- **Tectonic Renewal:** Repeated fault reactivation continuously exposes fresh surfaces for serpentinization while maintaining structural pathways, preventing system shutdown through complete reaction or sealing.

This creates a self-sustaining system where ancient materials remain chemically and structurally active rather than becoming passive fossil remnants.

Temporal Bridge to Mesozoic Integration & Assembly

The Mesozoic Farallon subduction event (180-50 Ma) represents the final major assembly phase that incorporated already-ancient materials into their current structural configuration, not their origin. Understanding this distinction is critical for recognizing how billion-year-old processes became integrated into California's modern geological framework.

Mulcahy et al. (2018) estimate that Franciscan subduction was occurring no later than 180 Ma, beginning in the Jurassic, and continuing to emplace material until about 25 Ma. The onset of Farallon plate subduction beneath western North America created the Franciscan subduction complex, consisting of deformed and metamorphosed sedimentary and volcanic rocks subducted to depths of 19-32 km (Wakabayashi, 2015).

Critically, oceanic lithosphere including serpentinized mantle fragments from earlier stages was subducted to depths of 30-60 km and metamorphosed under blueschist-facies conditions (>2 GPa, 200-500°C, 4-14°C/km geothermal gradient) before being tectonically incorporated into the accretionary wedge as exotic blocks. Blueschist occurs in scattered pods or belts in Franciscan Assemblage rocks, typically in association with serpentinite, with scattered exposures throughout the Diablo Range and southeastern Santa Cruz Mountains between Lexington Reservoir and Morgan Hill (Stoffer, 2002).

The formation and exhumation of these high-pressure assemblages represents a coordinated continental-scale tectonic event. Rocks that subducted to depths of up to 80 km all came up at the same time around 160 million years before present, with cooling rates up to ~140°C/Ma indicating extremely rapid exhumation (Rutte, 2020). Most significantly, both high- and low-T mafic blueschists are coeval and probably formed along 1000+ km of the North American Plate margin at ~160–155 Ma (Ukar, 2012).

The accretionary complex was tectonically assembled. Later Phanerozoic accretion (Farallon, Franciscan) represents piles of new material on top of older materials. The ancient, serpentinized mantle and crust were already at the continental edge, then came Paleozoic microplates and arcs, Mesozoic subduction of the Farallon Plate, and the Franciscan Complex, all obducted on top of older mantle.

Around 30 Ma, changes in plate configurations gradually changed the entire geologic framework of California as subduction gave way to transform faulting, resulting in formation of the San Andreas Fault system (Stoffer, 2002). The transition to transform tectonics (50 Ma-present) created the current San Andreas fault system, which continues to rework and expose ancient materials.

It's like a bookshelf that got knocked over sideways and then cracked in half. So, now ancient mantle material is at the surface (serpentinite); opalite/silica are along faults and shears; the Franciscan complex is slab fragments tilted, rotated, shattered; and fluids are still moving through fractures. This progression demonstrates how continental margins evolve through complete Wilson cycles while preserving evidence of earlier tectonic episodes.

Beyond Alluvial Fans

After widespread uplift and erosion across Santa Clara Valley, the area began to subside and receive sediment around 1–1.5 million years ago, forming the modern Santa Clara Basin. Seismic-reflection data indicate that strata within this basin are generally flat-lying and parallel to the ground surface, with alluvial sections at most 400 meters thick overlying irregular Mesozoic basement or horizontal Miocene-Pliocene sedimentary sections (Langenheim, 2015).

Conventional geological descriptions of Santa Clara County rely heavily on the term "alluvial fans" to explain surface deposits (Plan Bay Area, EIR 3.8, June 2021), but this terminology obscures more than it reveals.

While "alluvial fans" accurately describes the distribution pattern (water deposited sediments in fan-shaped formations) it provides no insight into the composition, chemistry, or origin of these materials. Geologists use "alluvial fans" as intellectual shorthand, but it avoids analyzing what the deposits actually consist of or why they exhibit the patterns observed.

The observed patterns (systematic depth zonation, organized geographic distribution, sustained chemical signatures, and integrated creek system behavior) indicate active subsurface processes maintaining surface expressions.

Creek System Chemical Zonation

Historic reconstructions demonstrate systematic chemical zonation where creek systems transition from mountain sources through distinct hydrochemical zones before reaching the bay, indicating organized geochemical processes rather than random alluvial transport.

Academic reconstruction of historic Bay Area ecology and watersheds shows only one permanent creek in the Santa Clara County area that flowed from the Santa Cruz mountains to the bay, alongside the Guadalupe River. These

1800-1850 reconstructions show Saratoga Creek running from the hills in south Saratoga, then flowing past a marsh, and into Sanjan Creek, which then met Guadalupe River at the bay tidal marshes (Beller, E. et al, Nov. 2010). Saratoga Creek (now connected to San Tomas Acquinas Creek), was also a source of the "alluvial fans" (Cal. Dept. of Consv. 2002).



Historical Conditions, Circa 1850 (SFEI, 2010).

The 1850 ecology maps show the area where Saratoga Creek flows alongside the freshwater marsh, and where it merges with Sanjan Creek, to be in a "wet meadow." Once Sanjan Creek passes U.S.-101 as it flows to the bay, the 1850 reconstructions show it entering an area designated "alkali meadow" (Beller, E. et al, Nov. 2010).

Alkali meadows occurred as an ecotone in the low areas between wet meadow and south San Francisco Bay tidal marshlands. As Reed (1862) describes the alkaline area just east of San José, "it was a medium between the two, it was neither like the upland nor like the Salt marsh, but it partook of the character of both." It included both lands subject to extreme high tidal flooding and poorly drained non-tidal soils.

Serpentine Terrane Hydrology

San Tomas Aquino Creek displays characteristic serpentine terrane behavior that contradicts normal alluvial drainage patterns. Like Calabazas Creek, San Tomas Aquino Creek channel lost definition well before reaching the Bay. Researchers have described the "scatters" of the creek around Hamilton Avenue, and the "sink" of the creek around Williams Road, a mile further north (Beller, E. et al, Nov. 2010).



Creek origins at the foothills. (SFEI, 2018).

Disconnected distributary channels flowed north below the terminus of the San Tomas Aquino Creek, hydrologically connecting diffuse wet season flows to Sanjon Creek and even above its terminus, the channel of San Tomas Aquino Creek was notably shallow and poorly defined (Beller, E. et al, Nov. 2010).

One early map even shows two termini of the creek, both labeled "sink of creek". These descriptions illustrate the shallow, discontinuous nature of the channel, which presumably spread into a marsh or disappeared into coarse gravels. This interpretation is further corroborated by GLO survey notes, which describe the creek in this reach as "very shallow" and about 4m wide (Beller, E. et al, Nov. 2010).

This behavior reflects natural fracture systems through serpentinized basement rather than typical alluvial drainage. The serpentine in this area explains the creek's unusual behavior. Serpentine forms when oceanic crust gets altered by hydrothermal fluids and creates highly fractured, permeable bedrock with extensive underground flow networks. The creek isn't actually "losing" water; it's following natural fracture systems through the serpentine bedrock, which is why it "scatters," develops multiple "sinks," and loses its defined channel before reaching the bay.

The Hydrogeochemical Flow Path

The hydrogeochemical flow path operates through distinct zones:

- **Mountain Recharge**: Meteoric water infiltrates fractured serpentine in the Santa Cruz Mountains, initiating water-rock interaction.
- **Deep Circulation**: Water descends through fault-controlled fracture networks, undergoing progressive serpentinization reactions that increase pH, Mg, and

alkalinity while temperatures reach 200-315°C at depth.

- **Lateral Flow**: Chemically evolved water moves laterally through permeable fault zones toward the valley floor.
- **Surface Expression**: Deep fluids mix with shallow groundwater and emerge through springs, creek baseflow, and diffuse seepage, creating the systematic chemical zonation.

These 'sinks' and 'scatters' represent locations where surface water reenters deep circulation pathways, not instances of net water loss. This is actually a well-documented phenomenon in serpentine terranes, where streams routinely disappear into fractured bedrock and emerge elsewhere as springs. The shallow, poorly defined channels described in the historical accounts are classic serpentine hydrology. The weathered serpentine creates clay-rich soils that promote diffuse, marshy flow patterns rather than well-incised channels.

Systematic Chemical Signatures

The surface chemistry of Santa Clara County reflects organized geochemical processes rather than random alluvial deposition. Historical ecological mapping from 1850 documents "alkali meadows" where creek systems flow, supporting rare, specialized ecosystems including saltgrass, alkali milk vetch, and common tarweed adapted to alkaline conditions (Beller et al., 2010).

Groundwater monitoring records from the California EPA GAMA Information System show selected well data from the Santa Clara foothills near fault zones around Saratoga, Monte Sereno, and Shannon. Several wells in Sanborn County Park and El Sereno Open Space Preserve show geochemical signatures consistent with geothermal or deep fluid influence, including: alkaline waters (310–993 mg/L as $CaCO_3$), elevated iron (up to 12,000–24,100 μg/L), manganese (225–680 μg/L), calcium (~110 mg/L), TDS (1,020–1,090 mg/L), and specific conductivity (1,583–1,728 μS/cm). (CalEPA GAMA, accessed July 14, 2025).

These values suggest strongly mineralized, reducing groundwater, potentially influenced by fault-driven fluid circulation. The Santa Cruz Mountain foothills along this faulted valley floor may represent an active hydrogeologic interface — a modern echo of a primordial soup.

Analysis of valley sediments shows that "most of the gravel sampled from five wells in the Santa Clara Valley was derived from Mesozoic rocks" with "abundant Franciscan greenstone derived from the southwest" (Anderson et al., 2016; Wentworth, 2005). These are systematic transport of specific ancient crustal materials.

Santa Clara County soils represent the surface expression of active ancient crustal systems rather than passive alluvial accumulation. The systematic chemical zonation, organized depth distribution of metamorphic minerals, and persistent alkaline surface chemistry indicate ongoing processes connecting deep serpentine-dominated systems with surface environments. Wells encountering serpentinite and greenstone at depths like 431m beneath "normal" sedimentary sequences confirm that the source systems producing these unique soil signatures remain structurally active and accessible (Stanley et al., 2002).

Rather than simple water-deposited fans, Santa Clara County preserves complex deposits that record continuous operation of integrated deep crustal systems spanning billions of years. The "alluvial" materials serve as carriers for ancient geochemical signatures, creating surface soil conditions that reflect deep-time geological processes still operating beneath the urbanized landscape.

Ongoing Deep-Time Processes

Multiple lines of evidence suggest Santa Clara County serpentinites preserve ancient mantle signatures and complex, multi-stage histories. Rather than representing straightforward oceanic crust fragments, these serpentinites exhibit geochemical, structural, and geophysical anomalies indicating deep mantle sources, multiple serpentinization episodes spanning significant time periods, and ongoing deep-seated processes that are difficult to reconcile with standard ophiolite models.

The evidence comes from isotopic signatures revealing ancient mantle melting events, geophysical anomalies suggesting deep-rooted serpentinite bodies, and structural relationships indicating complex, multi-stage alteration histories. Most significantly, several serpentinite bodies appear to represent active diapiric rise from forearc mantle wedge depths rather than passive ophiolite emplacement, with ongoing serpentinization processes at temperatures and pressures indicating deeper sources than typical ophiolite weathering.

The presence of extensive serpentine deposits on land today provides a geological timeline that, when traced backwards, points strongly toward Archean formation. Working backwards from serpentine currently exposed on continental surfaces, we can establish that most ophiolite emplacement events (where oceanic crust gets thrust onto land) occurred during major Precambrian orogenic episodes, typically between 800 million and 1.6 billion years ago. However, the oceanic crust involved in these collisional events was already geologically mature, usually requiring an additional 200-500 million years of seafloor existence before emplacement. This pushes the initial formation of the oceanic substrate back to approximately 1.0-1.6 billion years ago.

The serpentinization process itself requires substantial time to produce the extensive deposits we observe today. While modern examples like Lost City demonstrate serpentinization over tens of thousands of years for localized areas, regional-scale serpentinization sufficient to create large land-based deposits would require hundreds of millions of years of sustained hydrothermal activity. Adding this serpentinization duration to the oceanic crust formation timeframe pushes the initiation of serpentinization back to 1.5-2.0 billion years ago or earlier.

When we factor in the slower plate tectonics of early Earth which allowed oceanic crust to persist longer and undergo more extensive alteration, the timeline extends further backward, reaching past two billion years ago. This reconstruction, places the formation of today's land-based serpentine deposits squarely within the Archean Eon, demonstrating that temporal constraints alone are sufficient to establish this ancient origin.

The most likely time for the original serpentinization of San Francisco Bay Area mantle material was between 2.5 and 2.0 billion years ago, during the Late Archean to Paleoproterozoic periods.

This timing makes geological sense for several reasons. During this ancient era, mantle peridotite was accessible at or near the Earth's surface, the oceans had stabilized, and early subduction systems were operating. Crucially, these conditions allowed serpentinization to occur and to be preserved along the edges of ancient continents.

The serpentine we see today formed during this primordial window, when geological conditions were optimal for both the chemical transformation and long-term preservation of these materials. Subsequent tectonic events over billions of years then transported this already-serpentinized material to its current location along Laurentia's western margin.

Continuous operation over geological time scales requires specific preservation conditions:

- **Structural Isolation**: Serpentinized materials at continental margins avoid the high-temperature metamorphic events that would dehydrate and destroy serpentine in cratonic interiors.
- **Self-Sustaining Chemistry**: Serpentinization is exothermic and creates its own favorable conditions—released heat drives circulation, volume expansion maintains fracture permeability, and alkaline chemistry prevents mineral dissolution.
- **Tectonic Renewal**: Each major tectonic event (supercontinent assembly/breakup, subduction, transform faulting) reactivates and extends the fracture network without destroying the system, maintaining access to unreacted materials.
- **Episodic Reactivation**: Rather than continuous uniform activity, the system operates through punctuated episodes of high activity (during major tectonic events) alternating with lower-level background processes, enabling billion-year survival through multiple Wilson cycles.

In the Precambrian, the planet was different. The mantle was hotter which meant slower convection and fewer active subduction zones. Larger, longer-lived oceanic plates may have persisted for hundreds of millions to over a billion years. No modern-style plate tectonics occurred until ~1.0–1.5 Ga by many models. So, ancient oceanic lithosphere (including ultramafic mantle rock that became serpentinite) may have formed over 1 billion years ago and persisted without subducting until caught in later collisions.

In an ophiolite sequence the pillow basalts might be 1.5 Ga old (when the oceanic crust formed), but the ultramafic rocks (and thus the serpentine derived from them) could represent much older mantle material (potentially two+ Ga or Archean). It only got exposed at the seafloor through tectonics/faulting at 1.5 Ga and then underwent serpentinization. The serpentine actually represents the alteration of ancient mantle material.

Modern Expression of Ancient Framework

The evidence demonstrates that Santa Clara County represents the surface expression of active ancient crustal systems rather than passive preservation of fossil fragments. The systematic patterns observed today reflect ongoing operation of integrated deep crustal systems spanning billions of years, making this region a unique natural laboratory where deep-time geological processes continue to influence modern surface environments.

Modern geochemical conditions in Santa Clara County match optimal parameters for active serpentinization, demonstrating these ancient systems continue operating. Studies show serpentinization occurs optimally when temperature is between 200 and 315°C and when fluids are carbonate undersaturated (Klein et al., 2013). These conditions match thermal maturity indicators requiring generation at 2,100-2,700 meters depth documented in Santa Clara Valley oil samples, indicating ongoing deep thermal processes.

The mechanism connecting two Ga materials to modern surface conditions operates through three integrated steps:

- **Structural Access:** Active fault systems create fracture networks extending from surface to serpentinized basement at 8+ km depth, maintaining open pathways for fluid circulation.
- **Chemical Processing:** Groundwater percolating through these fractures reacts with ancient serpentinized materials under optimal temperature conditions (200-315°C), acquiring alkaline chemistry and elevated Mg, Cr, Ni concentrations.
- **Surface Expression:** This chemically evolved groundwater reaches surface through springs, creek baseflow, and diffuse seepage, creating the systematic soil chemistry that drove historical agricultural excellence and alkali meadow distribution.

The fault-controlled circulation ensures continuous renewal of reactive surfaces and sustained geochemical signatures.

The southwestern boundary of Santa Clara Valley is controlled by a complex zone of northeastward-vergent thrust and reverse faults collectively termed the "range-front fault system," including the Monte Vista, Berrocal, Shannon, and Sargent Faults (Stanley et al., 2002). These faults have accommodated approximately 3 km of cumulative shortening and 3 km of uplift since 3 Ma, with ongoing uplift rates of 0.8 mm/yr since 4.6 Ma based on apatite fission-track ages. (Stanley et al., 2002).

The curst in the SF Bay Area is thought to be about 20-23km thick from surface down to mantle, compared to ~28km under southern Santa Clara county and ~40km under the Sierra Nevada. (Healy, 1963; Eaton, 1963).

This active fault system reaches depths of 8 km or greater, where faults may steepen and merge with the San Andreas Fault system, creating deep structural connections that enable fluid circulation from mantle depths to surface. The range-front fault system functions as both a structural boundary and a hydro-geochemical gateway, continuously bringing ancient materials to the surface while establishing fracture networks that guide deep circulation systems.

The basin architecture shows complex multi-level structure with three distinct sedimentary units: the surface Santa Clara Basin plus at least two concealed deeper basins (Cupertino and Evergreen) at 2.5-3 kilometers depth (Langenheim, 2015).

Seismic velocity modeling reveals "substantial variations in basement rock properties" inconsistent with uniform sedimentary fill over stable basement, while gravity profiles suggest basement configurations controlled by ancient structural boundaries rather than simple depositional processes (Langenheim 2015).

Mercury mineralization throughout Santa Clara County provides direct evidence that multiple deep circulation systems operate as an integrated network. The New Almaden Mining District produced mercury from cinnabar (HgS) that forms as a hydrothermal mineral precipitating from ascending hot waters at shallow depths.

The formation mechanism demonstrates direct integration between serpentine alteration and hydrothermal circulation. Silica-carbonate type mercury deposits are associated with serpentinite altered to an assemblage of silica and carbonate minerals, with initial alteration consisting of addition of carbonate to serpentinite followed by introduction of silica into the central core (Rytuba et al., 2016). These deposits are localized along regional faults that separate serpentine from other rock, where serpentinite's relative impermeability serves to localize carbon dioxide-rich fluids from which mercury deposits formed.

This integration demonstrates that conventional geological mapping, when combined with deep-time analysis, reveals Santa Clara County as a unique natural laboratory preserving direct connections to Precambrian supercontinent processes within a well-characterized modern geological framework. The fault systems constitute active machinery that has operated for billions of years to expose, fracture, and circulate fluids through ancient crustal materials, ensuring that 2+ billion-year-old processes remain active and accessible.

Rather than simply cutting through static rock formations, these faults create dynamic pathways connecting deep serpentine-dominated basement with surface water systems. This ongoing tectonic activity ensures that deep-time geological processes continue to influence modern surface environments through fault-controlled exposure and circulation systems, making Santa Clara County a unique location where ancient continental margin processes remain directly observable and scientifically accessible.

Active Serpentinization Conditions

Serpentine is a group of phyllosilicate minerals that forms through one of Earth's most fundamental geological processes—serpentinization. The serpentine group consists of common rock-forming hydrous magnesium iron phyllosilicate minerals with the general formula $(Mg,Fe)_3Si_2O_5(OH)_4$, resulting from the metamorphism of minerals contained in mafic to ultramafic rocks (Klein & Bach 2009). The three primary polymorphs of serpentine are antigorite, lizardite, and chrysotile, each with distinct structural arrangements despite sharing similar chemical compositions. (C.G.S., 2002).

This reaction is highly exothermic and produces molecular hydrogen as a byproduct when iron-bearing minerals are involved (Schulte et al. 2010). During serpentinization, large amounts of water are absorbed into the rock, increasing the volume by 30-40% and reducing density.

This process has operated throughout Earth's history, playing a crucial role in global geochemical cycles and potentially in the origin of life. Serpentinization is fundamentally a hydration and metamorphic transformation of ferromagnesian minerals, such as olivine and pyroxene, in mafic and ultramafic rocks to produce serpentinite (Klein & Bach 2009; Klein et al., 2013). Modern geochemical conditions in Santa Clara County match optimal parameters for active serpentinization, demonstrating these ancient systems continue operating.

The heat released by complete serpentinization of 1 kilogram of peridotite is 250,000 joules, enough to raise the temperature of 1 liter of water by 50 degrees (Bach et al., 2006). This highly exothermic reaction explains the ongoing thermal signatures observed throughout the region's groundwater and oil systems, demonstrating continuous energy release from active serpentinization processes.

The peridotite protoliths of Santa Clara County serpentinites represent fragments of ancient mantle lithosphere formed during Archean crustal differentiation. The surface exposure of ultramafic rocks represents the most direct connection to this deep-time heritage, preserving relict primary mantle minerals within serpentinized assemblages. Original peridotite formation occurred at extremely hot mantle temperatures with ultramafic composition representing partial melting of primitive mantle sources.

Detailed investigation of serpentinite mélange in nearby Redwood City reveals at least three stages of serpentinization of a clinopyroxene-bearing harzburgite protolith, with cores containing surviving peridotite minerals plus lizardite ± antigorite + brucite + magnetite domains surrounded by lizardite + magnetite rims (Uno and Kirby, 2019).

The largest, freshest polygonal serpentinite blocks contain cores comprised of surviving peridotite minerals plus lizardite (lz) ± antigorite (atg) + brucite (brc) + magnetite (mag) domains surrounded by a ~10 cm thick peripheral rim of lizardite + magnetite. Stage-A serpentinization reactions are internally balanced except for addition of H_2O, Cs and Rb, while Stage-B serpentinization was open system with CaO, Cs and Rb migrating outside the blocks with addition of H_2O, U, Pb ±LREE.



Geologic setting of Santa Clara Valley showing subsurface extent of "major serpentine sheet," based on aeromagnetic anomaly. (Roberts and Jachens, 2003; Wentworth, 2010).

Several classes of mineral-filled veins were found in both cores and rims and indicate fluid pressures approaching lithostatic pressures that enabled these opening-mode fractures to occur. Such high-pressure hydrothermal conditions were important in promoting the serpentinization reactions as well as enabling a weak rheology consistent with a cold-intrusion/diapiric emplacement mechanism (Uno and Kirby, 2019). This complex internal structure records multiple episodes of fluid-rock interaction spanning significant time periods rather than single-stage alteration.

Modern Geochemical Evidence

Chemical analyses from active alteration sites demonstrate ongoing serpentinization processes throughout Santa Clara County. Samples F-72, F-73, and F-74 from underground workings at New Almaden mine represent "completely serpentinized lherzolite, unsheared and completely fresh" with compositions showing active hydrothermal alteration (Faust and Fahey, 1962). These samples contain 35.98-37.36% SiO_2, 36.99-38.54% MgO, and 13.50-15.00% H_2O, indicating substantial water incorporation characteristic of active serpentinization.

Depth-dependent chemical variations throughout the Santa Clara Valley groundwater system indicate progressive evolution through serpentine-dominated basement rather than simple meteoric recharge. The typical processes of chemical evolution of groundwater in coastal aquifers include cation exchange, carbonate precipitation, sulfate reduction, silicate weathering, clay precipitation, base exchange, and mixing. (Hanson, 2015).



Map of Loma Prieta earthquake study areas. "SCZ" is Santa Clara zone. (Schmidt, 2014).

Groundwater from depth-specific monitoring-well samples as well as depth-dependent and composite samples from water-supply wells shows a progressive evolution that includes cation exchange, carbonate precipitation, and sulfate reduction as water flows from the edges to the center of the basin (Hanson, 2015).

Most of the deeper monitoring and water-supply wells from the central part of the valley appear to have different major-ion chemistry and may represent different water from the deeper aquifers that does not receive recent streamflow or artificial recharge. In particular, some of the samples from the Eleanor Park site in Palo Alto are more saline and may represent very different waters, relative to the other deep monitoring wells. (Hanson, 2015).

Boron isotopes from monitoring and supply wells indicate a clear partition between the shallow waters that are related to artificial recharge and the older water from the lower-aquifer system. The boron isotopic signature from the ELNR monitoring wells is heavier than seawater, which is typical of adsorption of lighter boron from seawater over long periods of time. This groundwater is analogous to the old seawater sampled in aquifers of Monterey Bay (Hanson, 2003).

The boron isotope signature and the ages of this groundwater from the ELNR wells in Palo Alto indicate these waters may represent older diluted seawater that was never flushed out of the deeper aquifers along the southwestern margin of the San Francisco Bay (Hanson, 2015).

There is a wide range of groundwater ages. Carbon-14 data indicates that the groundwater in the upper 152m generally is less than 2000 yr old, and groundwater in the deeper aquifer layers generally ranges from 16,700 to 39,900 yr old. In addition, many of the wells in the center of the basin with deeper screens do not contain tritium. This would suggest that the water pumped from these wells does not contain recent recharge (Hanson, 2015).

Serpentinous bedrock and related overlying sediments are present in parts of the lower aquifer system along the west side of the valley in the Oak Hill/Los Gatos region (Oze et al., 2003; Newhouse et al., 2004). Analysis of samples from the WLLO monitoring site indicate that there were two episodes of deposition of serpentinous sediments that are enriched in chromium, and related trace metals, manganese, and nickel. These sediments occur at depths of ~227 and 239 m below land surface (Oze et al., 2003).

The confined aquifer systems show anomalous pressure responses, with pore pressure fluctuations considerably larger than expected for normal sedimentary aquifers, indicating connection to deeper circulation systems. Analysis of head levels within different stratigraphic units suggest that pore pressure fluctuations within the confined aquifer can be considerably larger than that observed in the unconfined water table (Carroll, 1991). (Schmidt and Burgmann, 2003).

The area where the historic Saratoga Creek used to run, pooling in a lake, and connecting to Sanjon (and where the creek may still run underground), is an area of Santa Clara with known serpentine and that included marshes and alkali meadows downstream before meeting the bay and Pacific Ocean. (SFEI, 2010). This area is also near the assumed concealed faults and unexplained magnetic anomalies, where unexplained damage occurred during the 1989 earthquake (Chase, 1992; Schmidt et al., 2014). This area also includes "major serpentine sheets" based on aeromagnetic anomaly data. (Roberts and Jachens, 2003; Wentworth, 2010).

Currently, this area resides between Lawrence Expressway and San Tomas Expressway (W-E), and Bayshore Freeway/U.S. 101 and Central Expressway(N-S). In 1850, this area was marked a vegetated wetland. (SFEI, 2010). Today the area is a mixture of residential, commercial, and industrial zoning. There was one recent large development project in this area that provided some modern deep-soil sampling.

This large development project (33.4 acres) occurred 15m west of San Tomas Aquino Creek and sits atop the ancient Saratoga Creek, which may still run underground. (NETR Online, 2014). The hydrology study report is five pages, and the geotechnical report is also brief and vaguely claims that Saratoga Creek was "filled" in 1968-1980. (Santa Clara, Santa Clara Square CEQA, 2015).

Project reports note the elevation of groundwater at the site fluctuates in relation to flow from San Tomas Aquino Creek and also notes a confined aquifer containing groundwater under positive pressure around 10.7m below the surface under four wells stretching from Coranado to Octavius Drive, and from Scott Blvd to Augustine Drive). Borings in this area reported variations of wet, grey sand/clay sediments with trace organics between 3-14m depth. (Santa Clara, Santa Clara Square CEQA, 2015).

Shallow 1.5m bore samples in this area also repeatedly reported dark grey gravels with "green residue" and "green staining." The green residue and green staining notes occurred at multiple sites round the property, reducing the odds it was local contamination. Borings for this project also repeatedly reported wet "olive-grey" sand and clay around 22-35ft, and wet olive, olive-grey, or olive-brown sand/clay around 4-9m deep. (Santa Clara, Santa Clara Square CEQA, 2015).

Several separate borings also reported "red sand" around this depth. In addition, a 4-4.5m segment was noted as "very moist – boring may fill with water by tomorrow." (Cal. EPA DTSC, Santa Clara Square Apartments, 2015).

During construction, soil was excavated and moved or removed, and 22,830,000 kg of soil was taken to a landfill, and 9,710 cubic meters were encased onsite in containment cells. Soil excavation generally ranged 7-9m below surface. (Cal. EPA DTSC, Santa Clara Square Apartments, 2016).

Groundwater testing in this area frequently found elevated levels of selenium (6-19 µg/L) (Cal. EPA DTSC, Santa Clara Square Apartments, 2015). Selenium is notable as it's a rare mineral on land but is common in ocean water and often found in serpentinite and pyrite, signaling evaporative or redox-driven mobilization from weathered bedrock.

In addition, vanadium was found in soil and groundwater. Vanadium is notable as it's a rare mineral generally associated with volcanic activity and former-marine hydrothermal terranes. Its presence in both soil (138 mg/kg) and groundwater (19 µg/L) implies the site as geochemical reaction with an active redox environment, alkaline pH, and dissolving iron oxides. Se and V together suggest persistent oxidative mobilization from sulfide- or oxide-bound sources. Further, the addition of solvent pollution in the area creates reducing conditions and metal leaching.

These studies also identified elevated levels of other substances including Ba, Cr, Co, Cu, Hg, Ni, and Zn. There were also high levels of magnesium in the soil up to 24,900 mg/kg, am amount typical of enriched serpentine soils. (Santa Clara, Santa Clara Square CEQA, 2015; Cal. EPA DTSC, Santa Clara Square Apartments, 2015).



Santa Clara circa 1850, excerpt for the area of the development analyzed showing Sanjon creek and alkali meadows. (SFEI, 2010).

This area intersects the historic Saratoga Creek and lays directly next to San Tomas Aquino Creek. The nearest mapped fault is Monte Vista-Shannon which is 12km southwest and where Saratoga and San Tomas Aquino Creeks enter the valley.

The Santa Clara Square development provides a modern subsurface snapshot that supports the framework while highlighting lost research opportunities. The systematic occurrence of green staining and olive sediment in multiple borings, combined with exceptionally high soil magnesium concentrations (24,900 mg/kg), indicates active interaction with serpentine-derived materials.

The elevated chromium, nickel, and cobalt levels represent the classic geochemical signature of ongoing serpentine weathering, while selenium and vanadium concentrations in both soil and groundwater suggest redox-active conditions characteristic of hydrothermal environments.

Most significantly, this development sits directly on the ancient Saratoga Creek bed (the only permanent creek that historically connected the Santa Cruz Mountains to the bay through documented alkali meadows) with strong evidence of historic hydrothermal activity.

Further, the confined aquifer under positive pressure (a natural spring) and notes about rapid groundwater infiltration suggest this ancient hydrological corridor remains active, whether through continued underground flow of Saratoga Creek itself or overflow from the rerouted San Tomas Aquino Creek, both of which originate from the same fault-controlled mountain sources.

Remarkably, these systematic geochemical anomalies were detected despite minimal investigation (five-page hydrology report) by consultants not specifically looking for deep-time geological signatures. The scale of contamination requiring removal (22.8 million kg) and the location at the intersection of ancient and modern creek systems made this area prime real estate for deep-time geological research, representing a rare accessible window into billion-year-old processes operating beneath Silicon Valley's urban landscape.

Active weathering of ancient mantle minerals demonstrates ongoing chemical processing of billion-year-old materials into modern surface environments. Weathering of ultramafic rocks and serpentinites in the Franciscan Complex produces serpentine soils containing high concentrations of Cr as well as other potentially toxic elements including Ni, Co, and Mn. Chromium concentrations in serpentine soils from Jasper Ridge Biological Preserve in the Central Coast Range reach 4,760 mg kg^{-1}, nearly three times greater than the serpentinite protolith (1800 mg kg^{-1}), demonstrating active concentration processes operating in modern surface conditions (Oze, 2004). The development project in Santa Clara on the ancient Saratoga Creek also reported elevated Cr, Ni, and Co in the soil and groundwater. (Cal. EPA DTSC, Santa Clara Square Apartments, 2016).

The diversity of chromium-bearing ancient minerals actively contributing to ongoing chemical evolution includes chlorite (0.3 Cr wt. %), enstatite (0.4 Cr wt. %), augite (0.7 Cr wt. %), chromite (10.8 Cr wt. %), magnetite (8.2-10.3 Cr wt. %), and an ultra fine-grained mixture of spinel and a silicate phase containing 13.3 Cr weight percent. Chemical analyses and X-ray microprobe maps demonstrate that Cr-spinels in these soils undergo incongruent dissolution progressively enriching the spinel toward a Cr-enriched end-member ($FeCr_2O_4$), indicating active mineral transformation processes rather than static preservation (Oze, 2004).

Analysis of valley sediments reveals abundant metavolcanic rocks and blue amphibole throughout the stratigraphic sequence, with blue amphibole concentrations increasing with depth to >17% at 304 meters (Anderson et al., 2016). Blue amphibole represents blueschist facies minerals formed under high-pressure metamorphic conditions, indicating that "alluvial" sediments derive from ongoing erosion of active high-pressure systems. (Hanson, 2015).

Depth-dependent variations in concentrations of lithium, boron, barium, fluoride, strontium, and iron reflect ongoing changes in sedimentary mineralogy and chemistry through continued water-rock interaction (Hanson, 2015).

Hydrothermal System Integration

Mercury mineralization throughout Santa Clara County provides direct evidence that multiple deep circulation systems operate as an integrated network rather than isolated processes. The systematic distribution of mineral deposits, hot springs, and hydrocarbon seeps reveals a unified deep system that has operated continuously for millions of years, connecting ancient serpentine-dominated basement with surface hydrothermal activity.

Mercury Mineralization: Tracing Deep Fluid Pathways

The New Almaden Mining District was the largest producer of mercury in North America. However, cinnabar (HgS) deposits were also discovered throughout Santa Clara County, including the Guadalupe Creek/Hicks Road area, Los Gatos Creek, and Deep Gulch (Bailey & Everhart, 1964; Santa Clara Co., 2012; Cal. Journal of Mines & Geology, 1949).

The formation mechanism demonstrates direct integration between serpentine alteration and hydrothermal circulation. Silica-carbonate type mercury deposits are associated with serpentinite altered to an assemblage of silica and carbonate minerals. The process involves initial alteration consisting of addition of carbonate to serpentinite followed by introduction of silica into the central core (Rytuba et al., 2016).

These deposits are localized along regional faults that separate serpentine from other rock, where serpentinite localizes carbon dioxide-rich fluids from which mercury deposits form. The deposits commonly extend up to several kilometers along fault zones, demonstrating that this is a regional-scale integrated system (U.S. EPA, 2005; Rytuba, 2003).

Mercury mineralization occurred at depths ranging from the surface to about 793m and at temperatures of 150°C, around the optimal range for serpentinization processes (Bailey and Everhart, 1964). The richest ore bodies were localized along gently dipping contacts where hydrothermal solutions rising from deep-seated sources spread out and stagnated under relatively impervious sheared rocks, demonstrating systematic fluid flow through organized pathways controlled by structural architecture.

Historical Mineral Springs: Century-Scale Evidence

Pacific Congress Springs and Regional Network

Multiple mineral springs throughout Santa Clara County document sustained deep hydrothermal circulation spanning over a century of commercial operations. Pacific Congress Springs near Saratoga contained 335.8 grains of dissolved minerals per gallon, including chloride of sodium, sulfate of soda, and carbonate compounds (Munro-Fraser, 1881). This represents highly mineralized water emerging from deep circulation through serpentine-rich basement.

Gilroy Hot Springs Complex

Gilroy Hot Springs maintained consistent 118°F temperatures with dissolved sulfur, iron, soda, magnesia, and trace arsenic, supporting commercial resort operations from 1866-1940s (Munro-Fraser, 1881; Santa Clara Co., 2012; Sanders, 2012). The springs were discovered in 1865 in the foothills northeast of Gilroy. In 1866, they were developed as the Gilroy Hot Springs Resort, which grew to include a hotel, cottages, bathing pools and bathhouses. The facilities expanded with Madrone Soda Springs, located 9.7km north of Gilroy Hot Springs. (Salewske 1982).

Systematic Regional Distribution

Similar sources of mineral water were exploited throughout the region during the late nineteenth century in Alum Rock Canyon and above Saratoga. Alum Rock became a regional park under San José management, and Saratoga Springs (also known as Congress Springs) was developed as a private park (Santa Clara Co., 2012). Soda Springs Creek and Soda Springs Road derive their names from a mineral spring located a short distance up the creek (Young 1984).

At least three regions of Santa Clara County are formally designated as hydrothermal systems: "White Sulphur Springs" (East Fork Coyote Creek), "Gilroy Hot Springs," and "Sargent Estate Warm Spring." White Sulphur Springs, closest to central Santa Clara Valley, reported 2,150 total dissolved solids per liter at 29°C (Cal. Dept. of Conserv. 1980).



Lines to depth of groundwater with area of "artesian flow" in Santa Clara Valley wells circa 1938 (Gardner, 1958).

Santa Clara County's artesian water discovery in the mid-1800s was considered transformational for the region (Sanders, 2012). Contemporary accounts described wells providing "pure, sweet, wholesome water" that emerged naturally from subterranean depths, creating what one historian called "wealth for this beautiful valley far beyond the value of a dozen gold mines" (Munro-Fraser, 1881). The artesian systems supported extensive agricultural development and were recognized as accessing deep groundwater circulation throughout the valley floor. Eventually, increased groundwater pumping lowered water levels, dropping water tables from near-surface to several hundred feet, and artesian wells became "*a thing of the past*" (USDA, 2015).

Integration with Hydrocarbon Systems

Unified Deep Circulation Networks

Integration with hydrocarbon systems confirms unified deep circulation networks operating through the same structural architecture that controls serpentinization and mercury mineralization. The Santa Clara Valley is underlain by a concealed, oil-bearing sedimentary basin that in some locations can be more than 3km thick, assumed to be created by an episode of normal faulting, volcanism, and crustal extension (Stanley, et al, 2002).

A conspicuous isostatic-gravity low extends about 35 km from Palo Alto southeastward to near Los Gatos, reflecting an asymmetric, northwest-trending sedimentary basin comprising low-density strata. The thickest accumulation (approximately 4 km thick) occurs along the basin's steep southwestern margin, which may be controlled by buried, northeast-dipping faults (Stanley, et al, 2002). The Evergreen low suggests a graben exists within the lower crust under Santa Clara Valley, which "may extend into the upper mantle" (Robbins, 1976).

Systematic Hydrocarbon Distribution

Oil near Los Gatos contained biomarkers typical of the Monterey Formation with stable carbon isotopic composition ($\delta 13C$ = -23.32‰) and thermal maturity indicators requiring generation at 2,100-2,700 meters depth (Stanley et al., 2002). Reports of oil around Moody Gulch and Saint Joseph's Hill in Los Gatos suggest an estimated 1 billion barrels of oil may have been generated in the deep subbasin between Los Gatos and Cupertino (Stanley et al., 2002).

Oil near New Almaden, often co-occurring with mercury mineralization, showed higher thermal maturity equivalent to 3,660m depth with distinctive biomarker patterns and evidence of thermal alteration by hydrothermal fluids (Stanley et al., 2002). At several localities in the New Almaden area, silica-carbonate rocks contain hydrocarbons that occur as droplets and films of tar and oil (Stanley et al., 2002).

Shared Fault-Controlled Pathways

The structural and chemical connections operate through shared fault-controlled pathways:

- **Structural Integration**: The range-front fault system creates deep fracture networks that tap multiple source regions; serpentinized basement (mercury + alkaline fluids), buried organic-rich sediments (hydrocarbons), and deep hydrothermal circulation (hot springs).

- **Chemical Integration**: Mercury precipitation requires temperatures <200°C, oil generation occurs at 100-150°C, and optimal serpentinization operates at 200-315°C (overlapping temperature windows that enable co-occurrence in the same fault-controlled system).
- **Depth Zonation**: Deep sources (serpentinization + thermogenic oil generation) feed upward through fault networks, with progressive cooling creating systematic precipitation/concentration zones—mercury closest to surface (<200°C), mixed hydrocarbon phases at intermediate depths, and deep serpentinization reactions at the base.

Regional Hydrocarbon and Sulfur Systems

Reports of sulfurous gas throughout the region provide additional evidence. Some wells penetrated intervals described as containing "brown sulfur" (90m) and "sulfur sand strong" (180m) (Stanley et al., 2002). Sargent's Oil Field produced over one million barrels of oil and at least 11.5 cubic meters of natural gas, with gas production attributed to thermogenic origin (Wagner, 2002). Multiple seeps were reported as "tar and sulphur water" (Hodgson, 1987).

Sixty acres of tar springs (oil and asphalt seeps) were reported in southern Santa Clara County around La Brea Creek and Sargent's Station, which was turned into a refinery (Santa Clara Co., 2012). Reports from Tar Creek at Sargent's included oil seeps "exuding from a serpentine foundation" and "canyon mouth seeps" within fifty feet of an "outcrop of heavy serpentine" (Wanger, 2002).

Studies hypothesized the deepest oil in the Sargent Oil Field Area "originated from a deeply buried, siliceous algal-rich source rock" deposited in a "marine, anoxic environment." The deepest oil sample was from 1,066m (Wanger, 2002).

Integrated Chemical Systems

Chemical, isotopic, and thermodynamic properties of CO_2-rich ground waters in the central California Coast Ranges demonstrate integrated metamorphic processes. The acidic CO_2-rich waters react with serpentine to form silica-carbonate rock, the host rock of many mercury deposits in the California Coast Range. The waters are partly of metamorphic origin and partly locally derived meteoric waters, with CO_2 entirely derived from metamorphic reactions at depth.

Depending on the relative importance of several reactions, the relative abundances of silica and carbonate minerals vary in the silica-carbonate rock. If CO_2-rich fluids react directly with peridotite or dunite, massive magnesite deposits may form (Barnes, et al, 1973).

This co-occurrence demonstrates that the same deep fracture systems transport metallic fluids (mercury), organic fluids (oil/gas), and hydrothermal fluids (hot springs) from depth, indicating comprehensive integration of serpentine alteration, thermal processes, and mineral systems throughout the region.

In Santa Clara County, hydrocarbon generation appears to occur at depths and temperatures consistent with deep hydrothermal circulation systems. This supports the model of ongoing deep crustal processes driving hydrology and mineral systems throughout the region, creating a vertically and laterally integrated system where different fluid types share pathways while maintaining distinct chemical signatures.

This integration demonstrates that conventional geological mapping, when combined with deep-time analysis, reveals Santa Clara County as a unique natural laboratory preserving direct connections to Precambrian supercontinent processes within a well-characterized modern geological framework. The fault systems constitute active machinery that has operated for billions of years to expose, fracture, and circulate fluids through ancient crustal materials, ensuring that 2+ billion-year-old processes remain active and accessible.

Modern Mineralogical Expression of Deep-Time Processes

The ongoing operation of Santa Clara County's ancient geological systems finds compelling expression in a remarkable assemblage of rare and structurally anomalous minerals documented from the region's hydrothermal localities. These species provide direct evidence for the systematic operation of exotic geochemical processes that reflect the continued activity of billion-year-old crustal systems.

Santa Clara County has produced an extraordinary concentration of mineralogical discoveries that challenge conventional understanding of mineral formation processes. Between 1912 and 1986, investigations documented six rare mineral species from the county's hydrothermal localities, including type locality species such as santaclaraite, franciscanite, melanophlogite, gageite, voelckerite, and kempite (Gjøvik, June 9 2025). Each represents a crystallographic anomaly that reflects the specialized geochemical conditions created by the ongoing operation of integrated deep crustal systems.

Melanophlogite stands as perhaps the most remarkable example. It's a natural silica clathrate where SiO_2 frameworks form hollow polyhedral cages hosting guest molecules including methane, carbon dioxide, and sulfur species. The Mount Hamilton occurrence remains the only locality worldwide where non-cubic crystal forms have been observed, with water-clear crystals reaching exceptional sizes up to 5 mm displaying unique tetrahexahedral modifications and internal hopper structures (Gjøvik, June 9 2025). This structural complexity requires precise temperature, pressure, and fluid chemistry conditions that reflect systematic rather than random hydrothermal processes.

Santaclaraite presents another structural exception as a hydrated calcium-manganese pyroxenoid with five-tetrahedral silicate chains, where hydrogen plays three distinct structural roles through both hydroxyl groups and structural water. This is a complexity rarely encountered in natural chain silicates. Franciscanite adds a vanadium-rich sheet silicate featuring distorted Mn and V octahedra in hexagonal symmetry, while gageite exhibits rare octahedral pipe-framework structures existing in multiple polytypic forms that reflect systematic crystallization under controlled conditions (Gjøvik, June 9 2025).

The geochemical characteristics of these minerals directly support the deep-time framework developed through isotopic and structural analysis. The concentration of manganese-bearing species (santaclaraite, franciscanite, gageite, kempite) reflects the systematic mobilization of transition metals through serpentinization processes.

Melanophlogite's incorporation of organic guest molecules demonstrates the reducing environments created by interaction between hydrothermal fluids and carbonaceous sediments within the accretionary complex. The structural dependence on these organic components for lattice stability reflects the complex fluid-rock interactions characteristic of active hydrothermal systems (Gjøvik, June 9 2025).

The systematic occurrence of these minerals within altered serpentinite and silica-carbonate rock assemblages of the Franciscan Complex indicates that the same structural architecture controlling modern mercury mineralization and hydrocarbon systems also governs exotic mineral formation. This integration demonstrates unified deep circulation networks operating through shared fault-controlled pathways rather than isolated hydrothermal episodes.

The specific mineralogical associations documented in Santa Clara County suggest hydrothermal systems capable of mobilizing normally incompatible elements under precisely controlled conditions that require billion-year-scale crustal evolution to achieve.

This mineralogical evidence supports Santa Clara County as a region that is an active natural laboratory where deep-time geological processes continue producing exceptional results observable through modern analytical techniques. The ongoing formation of structurally anomalous minerals provides direct proof that ancient systems remain active and accessible rather than representing passive preservation of fossil assemblages.

Structural Evidence for Continuous Operation

The structural architecture of Santa Clara County demonstrates ongoing tectonic processes that continuously expose, fracture, and circulate fluids through ancient crustal materials. The southwestern boundary of Santa Clara Valley is controlled by a complex zone of northeastward-vergent thrust and reverse faults collectively termed the "range-front fault system," including the Monte Vista, Berrocal, Shannon, and Sargent Faults (Stanley et al., 2002).

These faults have accommodated approximately 3 km of cumulative shortening and 3 km of uplift since 3 Ma, with ongoing uplift rates of 0.8 mm/yr since 4.6 Ma based on apatite fission-track ages. This active fault system reaches depths of 8 km or greater, where faults may steepen and merge with the San Andreas Fault system, creating deep structural connections that enable fluid circulation from mantle depths to surface.

The range-front fault system functions as both a structural boundary and a hydro-geochemical gateway, continuously bringing ancient materials to the surface while establishing fracture networks that guide deep circulation systems. This active fault system creates a critical structural boundary in the Saratoga area where San Tomas Aquino and Saratoga Creeks originate (Stanley, et al, 2002).

This zone represents the transition from exposed serpentine and other sea crust/mantle complexes in the Santa Cruz Mountains to the concealed deep basin systems beneath the valley floor. The fault-controlled topography creates a natural structural vault where groundwater from the mountains encounters the buried serpentine basement, establishing the initial geochemical signatures that these creeks then transport northward through the valley floor.

The Silver Creek fault beneath Santa Clara Valley creates "a sharp deformation gradient in interferometric synthetic aperture radar (InSAR) data" that results from "partitioning of the basin aquifer by the fault" rather than tectonic slip alone (Langenheim, 2015). This demonstrates ongoing interaction between active faulting and deep groundwater circulation systems.

Current geodetic measurements show the Sargent Fault slipping at 3 mm/yr, while geodetic data indicates "northeast-directed contraction continued after the earthquake along the southwest margin of Santa Clara Valley" (Stanley et al., 2002; Schmidt et al., 2014). This ongoing deformation continuously creates new fracture networks and reactivates existing ones, maintaining pathways for deep fluid circulation.

Boron isotope studies reveal two distinct groups of serpentinites: those associated with blueschists show lighter boron isotope signatures ($\delta 11B$ = -12 to +8.8‰) interpreted as interaction with forearc slab fluids at deeper depths, while others show heavier signatures ($\delta 11B$ = +7.2 to +24.4‰) (Boschi et al. 2019; Kirby et al. 2019). This isotopic discrimination indicates multiple fluid sources and alteration environments spanning significant depth ranges, demonstrating ongoing fluid circulation through fault-controlled pathways that connect deep serpentine-dominated basement with surface water systems.

The 1989 Loma Prieta earthquake revealed systematic damage patterns that illuminate the hidden structural architecture controlling ongoing processes. In the Santa Clara zone, diffuse damage occurred 8 km northeast of the Santa Cruz Mountains over alluvium, following patterns aligned with aeromagnetic anomalies indicating concealed serpentinite bodies and gravity gradients marking basin margins (Schmidt et al., 2014). "Structural data for this area are sparse." (Cal. Dept. of Consv. 2002).

Near Los Gatos and Cupertino, concrete showed "remarkably regular patterns of deformation consisting of buckling and thrusting in a north-south to northeast-southwest direction," identical to damage patterns from the 1906 earthquake, indicating systematic structural control that operates repeatedly. (Schmidt et al., 2014).



Deformed storm-drain grate in Los Gatos on state Highway 17 at Route 9, following Loma Prieta 1989 earthquake. (Schmidt, 2014).

Coast Range serpentinites show evidence of multiple deformation episodes with different mineralogical assemblages and cross-cutting relationships, including in-situ mantle alteration, crustal-level hydration, and late-stage shearing, indicating complex burial-exhumation cycles rather than simple emplacement (Kirby et al. 2019). The fault systems constitute the active machinery that has operated for billions of years to expose, fracture, and circulate fluids through ancient crustal materials. The earthquake damage patterns reveal that this structural architecture extends throughout the urbanized valley, creating a hidden network of fault-controlled fluid pathways that connect the exposed ancient systems in the mountains with the concealed ancient systems beneath Santa Clara County.



Aeromagnetic Map of the San Francisco-San Jose Quadrangle, California (Chase, 1992).

This ongoing tectonic activity ensures that 2+ billion-year-old processes remain active and accessible, making Santa Clara County a unique location where deep-time geological processes continue to influence modern surface environments through fault-controlled exposure and circulation systems.

A Note on "Jurassic Rocks" and "Alluvial Fans"

The Scientific Information Vacuum

Santa Clara County represents one of the most economically significant and resource-rich regions on Earth. Silicon Valley hosts the world's leading technology companies, major research universities, and substantial government research facilities. Despite this concentration of scientific and financial resources, and the region's obvious geological importance, there is remarkably little comprehensive geological research.

While industry conducts private geological assessments for development and operational purposes, the publicly accessible geological literature for Santa Clara County remains extraordinarily limited. Standard geological references describe the region in terms of "alluvial fans" and

"Jurassic rocks." For a region of such scientific and economic importance, this represents an extraordinary gap in accessible geological knowledge.

Geological reports from the 1950s through 1980s exhibit consistent patterns of limited discussion regarding pre-Jurassic geological formations and deep-time processes, despite documented evidence for much older assemblages. This pattern coincided with several institutional pressures that influenced scientific communication during this period.

The legacy of the Scopes trial (1925) had established legal and cultural precedents affecting educational content related to deep geological time. California became a focal point for organized resistance to deep-time geological content, with institutions like the Institute for Creation Research (founded 1970) pursuing litigation strategies that specifically targeted textbook publishers and educational institutions, working to influence "government at the legislative and judicial levels" (Bergman, 1993).

Pre-Jurassic marine geology was particularly problematic for these groups because it documented hundreds of millions of years of marine life evolution prior to terrestrial ecosystems (Gjøvik, June 25 2025); evidence that directly contradicted specific religious narratives about Earth's history.

The scientific community's limitations prior to the acceptance of plate tectonic theory in the late 1960s provided additional justification for avoiding detailed temporal discussions of complex geological formations. By the time comprehensive explanatory frameworks became available, institutional patterns of limited deep-time discussion were already established.

The discovery of hydrothermal vent origins of life (1970s-2000s) increased institutional resistance to deep marine geological content, as this research suggested life originated in deep ocean environments through chemical processes. (Gjøvik, June 23 2025).

Religious institutional presence, including the active Catholic Mission Santa Clara de Asís within Santa Clara University and Santa Clara's designation as "The Mission City," may have also reinforced cultural pressures.

During the technology sector expansion of the 1990s-2000s, economic development interests may have found common ground with groups opposing comprehensive geological education. Detailed understanding of deep-time geological processes has significant implications for development costs, infrastructure planning, and land use creating economic incentives for simplified geological interpretations in public planning documents.

The Consequence: Lost Scientific Opportunity

The result of these converging pressures is that one of Earth's most remarkable geological laboratories has remained scientifically underexplored in publicly accessible literature. The evidence presented in this study suggests Santa Clara County preserves active geological systems spanning two billion years. This makes it potentially one of the most scientifically significant regions for understanding deep-time Earth processes globally.

This represents more than academic oversight. The systematic avoidance of comprehensive geological investigation in such a scientifically important region suggests institutional mechanisms that can effectively suppress critical scientific information when it conflicts with powerful religious and economic interests. (Scott, 2016).

The *Santa Clara County case* demonstrates how institutional pressures can create persistent gaps in scientific understanding, even in regions of obvious scientific importance. The convergence of religious opposition to deep-time evidence, economic development interests, and institutional inertia can apparently maintain scientific information gaps across multiple decades, even in the presence of abundant physical evidence.

This has broader implications for scientific practice, particularly regarding how institutional and cultural pressures influence which research questions are pursued, how results are communicated, and what information becomes accessible to the broader scientific community and public.

The academic geological community must recognize that conducting independent, comprehensive geological research may sometimes require addressing institutional factors that have influenced previous scientific communication, particularly when those factors may have impeded full investigation of scientifically significant phenomena.

Academic Responsibility

The evidence presented in this study suggests that Santa Clara County's geological systems may represent one of the most complete and accessible records of deep-time Earth processes available for scientific investigation. The decades-long gap in comprehensive geological analysis of this region highlights the importance of academic freedom in pursuing scientific investigation regardless of institutional sensitivities or economic considerations.

Future geological research must address acknowledged gaps in regional understanding through systematic application of modern analytical techniques across all relevant temporal frameworks, ensuring that scientific investigation proceeds according to evidence rather than institutional constraints that may have influenced previous research patterns.

Future Research

The deep-time framework presented here opens numerous avenues for targeted research that could further test, refine, and extend our understanding of Santa Clara County's ancient geological heritage. The following research directions represent high-priority investigations that could significantly advance both regional geological knowledge and broader understanding of how continental margins preserve and maintain billion-year-old crustal systems.

Structural and Geophysical Investigations

Expanded Geochronological Analysis: Systematic U-Pb zircon dating of additional Coast Range Ophiolite samples could confirm the extent of 1.5+ Ga signatures and map their distribution across the region. Combined Lu-Hf isotopic analysis would provide insights into the mantle source characteristics and crustal recycling processes during supercontinent cycles.

Re-Os Isotope Mapping: Detailed osmium isotope analysis of serpentinites across Santa Clara County could trace the spatial extent of ancient mantle domains and identify areas where Archean signatures might be preserved. This work could definitively establish whether two+ Ga materials are present and map their distribution.

Boron Isotope Systematics: Expanded boron isotope analysis of serpentinites and associated fluids could distinguish between different fluid sources (metamorphic vs. meteoric) and trace the evolution of fluid compositions through different alteration episodes, providing insight into the timing and character of ancient serpentinization events.

Deep Seismic Profiling: High-resolution seismic reflection profiles across the range-front fault system could image the deep structural architecture and confirm whether fault networks extend to mantle depths as predicted by the deep-time model.

Magnetotelluric Surveys: Detailed magnetotelluric studies could map the distribution of conductive serpentinized zones at depth and trace their connections to surface hydrothermal systems, testing predictions about integrated deep circulation networks.

Gravity and Magnetic Modeling: Advanced 3D modeling of gravity and magnetic anomalies could refine estimates of serpentinite body geometry and test whether the observed patterns are consistent with billion-year preservation rather than Mesozoic emplacement.

Geochemical and Hydrological Studies

Noble Gas Geochemistry: Analysis of helium, neon, and argon isotopes in hydrothermal fluids could distinguish between mantle-derived and crustal components, providing direct evidence for ongoing deep mantle contributions to surface hydrothermal systems.

Systematic Water Chemistry: Comprehensive sampling of springs, wells, and creek systems throughout Santa Clara County could map the full extent of deep fluid contributions and test predictions about systematic chemical zonation reflecting ancient basement control.

Trace Element Fingerprinting: Detailed analysis of trace element signatures in sediments, soils, and waters could trace contributions from specific ancient lithologies and test whether modern surface chemistry reflects ongoing interaction with billion-year-old materials.

Environmental Isotope Tracing: Use of stable isotope tracers to follow nutrient and water movement from deep sources through surface ecosystems could quantify the magnitude of deep geological influence on modern environmental systems.

Comparative Studies

Regional Context: Comparison with other ophiolite complexes worldwide could determine whether billion-year preservation is unique to Santa Clara County or represents a previously unrecognized phenomenon that might exist elsewhere along ancient continental margins.

Analog Studies: Investigation of similar geological settings in other regions could test whether the deep-time preservation model applies more broadly and identify other locations where ancient continental margin processes might remain active.

Archean Analogs: Comparison of Santa Clara County serpentinization systems with Archean hydrothermal deposits could test whether these modern systems provide insights into early Earth processes and environments.

Advanced Analytical Techniques: Application of emerging analytical methods (e.g., atom probe tomography, in-situ laser ablation techniques) could reveal micro-scale evidence for ancient processes preserved within individual mineral grains.

Conclusions

A Paradigm Shift in Regional Understanding

This study presents compelling evidence that Santa Clara County preserves far more than scattered fragments of

Mesozoic oceanic crust. Instead, the systematic integration of isotopic, geochemical, structural, and ecological evidence reveals active preservation of geological systems spanning two billion years: making this region a unique window into Earth's deep-time evolution operating within a modern urban environment.

The convergence of multiple independent lines of evidence supports this transformative interpretation. Recent isotopic studies document 1.5+ billion-year model ages directly linking local materials to Nuna supercontinent processes. Systematic surface patterns demonstrate ongoing operation of integrated deep systems rather than passive preservation of ancient fragments. Modern geochemical conditions match optimal parameters for active serpentinization, while fault-controlled circulation systems maintain continuous access to ancient materials through structural networks that have operated for billions of years.

Key Evidence for Active Deep-Time Systems

Coast Range Ophiolite peridotites preserve aluminachron model ages of ~1.5 Ga, with researchers explicitly noting potential connections to Columbia supercontinent rifting. Point Sal peridotites show extensive melt depletion with osmium isotope ratios indicating long-term preservation of refractory mantle domains. These signatures demonstrate preservation of billion-year-old mantle materials, not formation of new oceanic crust 160 million years ago.

Historical agricultural excellence across diverse crops spanning 200+ years, systematic earthquake damage patterns following concealed geological boundaries, and organized chemical zonation from mountain sources through alkali meadows all indicate active subsurface processes maintaining surface expressions. These patterns require sustained operation of integrated systems, not episodic deposition or passive remnants.

Mercury mineralization, hydrocarbon seeps, and long-term hydrothermal activity demonstrate unified deep circulation systems operating through shared fault-controlled pathways. The co-occurrence of metallic fluids, organic compounds, and mineral-rich waters at optimal serpentinization temperatures reveals comprehensive integration spanning multiple depth zones.

Active fault systems reaching mantle depths, ongoing uplift rates, and systematic damage patterns indicate that ancient materials remain structurally accessible and continue to influence modern surface processes through fault-controlled exposure and circulation.

Redefining Continental Margin Evolution

If correct, this interpretation fundamentally changes how we understand the preservation of ancient crustal systems. Rather than viewing ophiolites as isolated snapshots of ancient ocean basins, Santa Clara County demonstrates that continental margins can preserve and maintain active connections to deep-time processes spanning multiple Wilson cycles and supercontinent assemblies.

This has profound implications for understanding how Earth's crust evolves and preserves evidence of ancient processes. The fault systems constitute active machinery that has operated for billions of years, continuously exposing, fracturing, and circulating fluids through ancient materials while maintaining their chemical and structural integrity. This creates a self-sustaining system where deep-time processes remain observable and scientifically accessible within modern geological frameworks.

Windows into Early Earth Processes

Santa Clara County's preserved oceanic sequences and hydrothermal systems provide direct analogs for understanding early Earth processes that operated under different atmospheric and oceanic conditions. Ancient seafloor preservation parallels conditions that may have existed in Archean oceans, while hydrothermal systems preserve chemical signatures similar to those proposed for early life environments. High-pressure metamorphic processes record deep crustal conditions relevant to early continental formation.

The ongoing operation of serpentinization systems that may have initiated during Archean time offers unprecedented opportunities to study how these fundamental Earth processes have evolved and adapted through changing global conditions while maintaining their essential character.

Natural Laboratory for Deep-Time Research

Perhaps most significantly, Santa Clara County represents a unique natural laboratory where deep-time Earth processes remain directly observable and accessible for study. The integration of ancient materials with modern analytical capabilities creates opportunities to investigate billion-year-old systems using twenty-first-century science.

This accessibility is particularly valuable because most ancient geological materials have been either destroyed by subsequent geological processes or buried at depths inaccessible to direct study. Santa Clara County's preservation of active ancient system creates opportunities for sustained detailed investigation that could advance fundamental understanding of how Earth systems operate across deep time.

Beyond Conventional Timescales

This study demonstrates the necessity of expanding conventional geological thinking beyond typical timescales when encountering systematic patterns that resist explanation through standard models. The acknowledged failures of existing frameworks to explain Santa Clara County's geology point to the need for deep-time perspectives that can account for billion-year preservation and operation of integrated crustal systems.

The methodology employed here (systematic integration of isotopic, geochemical, ecological, and structural evidence) demonstrates the power of interdisciplinary approaches to geological problems. The convergence of independent lines of evidence provides much stronger support for conclusions than any single analytical approach could achieve. This integrated methodology could be applied to other geologically complex regions where conventional approaches have failed to provide satisfactory explanations for observed systematic patterns.

A Living Connection to Deep Time

Santa Clara County's preservation of active billion-year-old geological systems creates a living connection to Earth's deep-time evolution that extends far beyond academic interest. The ongoing operation of ancient processes demonstrates that deep time actively shapes modern environments and continues to influence contemporary geological, hydrological, and biological systems.

This connection between ancient and modern processes offers profound insights into how Earth systems maintain continuity across geological time while adapting to changing conditions. The fault-controlled circulation systems that maintain access to ancient materials represent a form of geological memory that preserves and transmits information about Earth's evolution across billions of years.

Understanding and protecting these connections becomes not just a matter of scientific interest, but of preserving unique windows into Earth's deep-time heritage for future generations of researchers and society as a whole.

The transformation of our understanding of Santa Clara County from a collection of chaotic geological fragments to an integrated natural laboratory preserving two billion years of Earth history opens new perspectives on how ancient processes continue to shape modern environments and how continental margins preserve and maintain connections to deep time that remain scientifically accessible and environmentally significant today.

Acknowledgements

The author conceptualized the study, developed the theoretical framework, and conducted all analyses. This research received no external funding and was conducted using personally acquired resources.

The author acknowledges Claude AI and Scholar GPT for their assistance with manuscript preparation, data interpretation discussions, and research development over the course of this investigation.

Data Availability: No new data was produced in the creation of this article.

References

Adam, D. Hardcore hibernation. *Nature* (2000).

Agard, Philippe & Yamato, Philippe & Jolivet, Laurent & Burov, Evgene. (2009). Exhumation of oceanic blueschists and eclogites in subduction zones: Timing and mechanisms. Earth-Science Reviews. 92. 53-79.

Altermann, Wladyslaw. (2001). The oldest fossils of Africa - A brief reappraisal of reports from the Archean. Journal of African Earth Sciences. 33. 427-436. 10.1016/S0899-5362(01)00089-6.

Anderson, D.W. et al, 2016. Composition of sediment records late Quaternary paleogeographic evolution of Santa Clara Valley, California. *Geosphere*; v. 12, no. 2

Andreani, M., et al. "Serpentinization-Driven H_2 Production From Continental Break-Up to Mid-Ocean Ridge Spreading: Unexpected High Rates at the West Iberia Margin." *Frontiers in Earth Science* 9 (2021): 673063.

Bailey, E.H., and Everhart, D.C., 1964, Geology and quicksilver deposits of the New Almaden District, Santa Clara County, California: U.S. Geological Survey Professional Paper 360.

Baranov, V., 1957, A new method for interpretation of aeromagnetic maps: Pseudo-gravimetric anomalies: *Geophysics*, v. 22, p. 359-383.

Barnard, Patrick & Schoellhamer, David & Jaffe, Bruce & Mckee, Lester. (2013). Sediment transport in the San Francisco Bay Coastal System: An overview. *Marine Geology*. 345. 3–17.

Barnes, Ivan, V. C. LaMarche Jr., and Glen Himmelberg. "Geochemical Evidence of Present-Day Serpentinization." *Science* 156, no. 3776 (1967): 830-832.

Barnes, I. et al, (1973), Silica-carbonate alteration of serpentine: Wall rock alteration in mercury deposits of the California Coast Ranges, *Economic Geology*.

Barth, Andrew & Tosdal, Richard & Wooden, Joseph & Howard, K. (1997). Triassic plutonism in southern California: Southward younging of arc initiation along a truncated continental margin. Tectonics. 16.

Barth, Andrew P., et al. "Assembling and disassembling California: A zircon and monazite geochronologic framework for Proterozoic crustal evolution in southern California." *The Journal of Geology* 117.3 (2009): 221-239.

Beller, E. Historical Vegetation and Drainage Pat Terns of Western Santa Clara Valley, Nov. 2010, San Francisco Estuary Institute.

Bergman, Jerry. A Brief History of the Modern American Creation Movement, *Contra Mundum* No. 7 Spring 1993.

Bishop, C.C., and Williams, J.W., 1974, Thickness of alluvium, Santa Clara County, California, *in* Rogers, T.H., and Williams, J.W., eds., Potential seismic hazards in Santa Clara County, California: California Division of Mines and Geology Special Report 107, 39 p.

Blake, M.C., Jr., Bailey, E.H., and Wentworth, C.M., 2016, The Cedars untramafic mass, Sonoma County, California (ver. 1.2, April 2016): U.S. Geological Survey Open-File Report, 2012–1164, 13 p., 1 plate.

Blake, M. C., et al, Geologic Map and Map Database of Western Sonoma, Northernmost Marin, and Southernmost Mendocino Counties, California, Miscellaneous Field Studies Map MF-2402, USGS (2002).

Blakely, R.J., 1995, Potential theory in gravity and magnetic applications: Cambridge University Press.

Blewett, G. et al., "Multiple episodes of serpentinite alteration revealed by progressive leaching experiments," *Geochimica et Cosmochimica Acta* 385 (2025): 86-103.

Bonnet, G. et al, 2024. Seamount Subduction Dynamics and Long-Term Evolution of the Franciscan Active Margin. Tectonics, 2024, 43 (3), pp.e2023TC008084.

Boschi, Chiara, et al. "Boron isotope variations of Franciscan serpentinites, northern California." *Lithos* 336-337 (2019): 142-157.

Bowen, Oliver E. (1962). Geologic Guide to the Gas and Oil Field of Northern California, California Division of Mines and Geology, Bulletin 181, San Francisco.

Brabb, E.E., and Hanna, W.F., 1981, Maps showing aeromagnetic anomalies, faults, earthquake epicenters, and igneous rocks in the southern San Francisco Bay region, California: U.S. Geological Survey Geophysical Investigations Map GP-932, 6 p.

California Department of Conservation, (1980). Geothermal Resources of California, Division of Mines and Geology, California Geologic Data Map Series No. 4.

California Department of Conservation, (2000), Abandoned Mine Lands Unit, California's Abandoned Mines A Report on the Magnitude and Scope of the Issue in the State Volume I, Office of Mine Reclamation, (June 2000).

California Department of Conservation, (2002), Seismic Hazard Zone Report For The San Jose West 7.5-Minute Quadrangle, Santa Clara County, California, Seismic Hazard Zone Report 058.

California Environmental Protection Agency, Statewide Advisory for Eating Fish from California Rivers, Streams, and Creeks without Site-Specific Advice, Office of Environmental Health Hazard Assessment (OEHHA), California EPA, September 21, 2022.

California Environmental Protection Agency, Water Boards, GAMA Groundwater Information System, https://www.waterboards.ca.gov/gama/online_tools.html

California Environmental Protection Agency, (2015-2025). DTSC Id. No. 60002212**,** Santa Clara Square Apartments.

California Geological Survey, Serpentine: California state Rock, Conservation, Note 14 (2002).

California Journal of Mines and Geology, (1949), State of California, Division of Mines, Volume 45, Issue 1.

California Journal of Mines and Geology, (1949), State of California, Division of Mines, Volume 45, Issue 3.

California Journal of Mines and Geology, (1949), State of California, Division of Mines, Volume 45, Issue 4.

Cannaò, Enrico, et al. "A continental forearc serpentinite diapir with deep origins: Elemental signatures of a mantle wedge protolith and slab-derived fluids at New Idria, California." *Lithos* 396-397 (2021): 106-124.

Cannat, M., D. Bideau, and R. Hébert. "Serpentinization: A Review." *Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary Science* 325, no. 2 (1997): 175-181.

Cao, X. et al, (2024). Earth's tectonic and plate boundary evolution over 1.8 billion years. *Geoscience Frontiers*, Volume 15, Issue 6, November 2024, 101922.

Carter, W.R., and Jachens, R.C., 1993, Aeromagnetic map of the San Jose 1:100,000 scale quadrangle, California: U.S. Geological Survey Open-File Report 93–277, 1 sheet.

Cawood, P.A. et al, Linking collisional and accretionary orogens during Rodinia assembly and breakup: Implications for models of supercontinent cycles. 2016. Earth and

Planetary Science Letters, Volume 449, p. 118-126.

Chase, Gordon W. (1992). Aeromagnetic Map of the San Francisco-San Jose Quadrangle, California, 1:250,000. California Regional Geological Map Series, Map 5D (Aeromagenetic Anomaly).

Cho, H. et al, (2025). Imaging the 3D basin structure of the Santa Clara Valley by ambient noise tomography, *Geosciences Journal* (2025) 29:87–101.

Choi, Sung Hi & Shervais, John & Mukasa, Samuel. (2008). Supra-subduction and abyssal mantle peridotites of the Coast Range ophiolite, California. *Contributions to Mineralogy and Petrology*. 156. 551-576.

Choi, Sung Hi & Mukasa, Samuel & Shervais, John & Puchtel, Igor. (2025). PGE Abundance And Re-Os Isotopic Systematics Of Coast Range Ophiolite Peridotites And Chromitite, And Insights Into Fore-Arc Magmatic Processes.

Choi, Sung Hi & Mukasa, Samuel & Shervais, John & Puchtel, Igor. (2024). Re–Os Isotope and PGE Abundance Systematics of Coast Range Ophiolite Peridotites and Chromitite, California: Insights into Fore-Arc Magmatic Processes. Lithosphere. 2024.

Chowdhury, Priyadarshi & Cawood, Peter & Mulder, Jacob. (2025). Subaerial Emergence of Continents on Archean Earth. Annual Review of Earth and Planetary Sciences. 53.

Christensen, Nikolas. (2004). Serpentinites, Peridotites, and Seismology. International Geology Review - INT GEOL REV. 46. 795-816.

Cisneros, Miguel & Behr, Whitney & Platt, John & Anczkiewicz, Robert. (2022). Quartz-in-garnet barometry constraints on formation pressures of eclogites from the Franciscan Complex, California. *Contributions to Mineralogy and Petrology.*

Clarke, Anthony & Kirkland, Christopher & Glorie, Stijn & Quentin de Gromard, Raphael & Tucker, Naomi. (2024). Detrital zircon and apatite reveal Paleoproterozoic rifting along the eastern margin of the Yilgarn Craton. Precambrian Research. 414.

Condie, K. C. and Stern, R. J. (2023). Ophiolites: Identification and tectonic significance in space and time, Geoscience Frontiers, Volume 14, Issue 6, November 2023.

Cruz-Uribe, Alicia & Feineman, Maureen. (2009). Detailed geochemical analysis of a garnetite vein in eclogite, Ring Mountain, CA. AGU Fall Meeting Abstracts.

Dalziel, I. W.D. "Neoproterozoic-Paleozoic geography and tectonics: Review, hypothesis, environmental speculation," *Geological Society of America Bulletin* 109, no. 1 (1997): 16-42.

Debret, Baptiste & Andreani, M. & Munoz, Manuel & Bolfan, Nathalie & Carlut, Julie & Christian, Nicollet & Schwartz, Stéphane & Trcera, Nicolas. (2014). Evolution of Fe redox state in serpentine during subduction. Earth and Planetary Science Letters. 400. 206–218.

Dellinger, D. A. California's Unique Geologic History and Its Role in Mineral Formation, with Emphasis on the Mineral Resources of the California Desert Region. U.S. Geological Survey Circular 1024. (1989).

Dibblee, T.W., Jr., and Brabb, E.E., 1978, Preliminary geologic map of the Los Gatos Quadrangle, Santa Clara and Santa Cruz Counties, California: U.S. Geological Survey Open-File Report 78-453, scale 1:24,000.

Dilek, Yildirim. "Tectonic significance of post-accretion rifting of a Mesozoic island-arc terrane in the northern Sierra Nevada, California." *The Journal of Geology* 97, no. 4 (1989): 503-518.

Dilek, Yildirim. "Structure and tectonics of an Early Mesozoic oceanic basement in the northern Sierra Nevada metamorphic belt, California: Evidence for transform faulting and ensimatic arc evolution." *Tectonics* 8, no. 5 (1989): 999-1014.

Dilek, Yildirim, Peter Thy, and Eldridge M. Moores. "Episodic dike intrusions in the northwestern Sierra Nevada, California: Implications for multistage evolution of a Jurassic arc terrane." *Geology* 19, no. 2 (1991): 180-184.

Dickinson, W. et al. "Alternate origins of the Coast Range ophiolite (California): Introduction and implications." *GSA today* 6, no. 2 (1996): 1-10.

Dilek, Yildirim. (2003). Ophiolite concept and its evolution. *Special Paper of the Geological Society of America*. 373.

Dilek, Y. and Newcomb, S., Ophiolite concept and the evolution of geological thought, *Geological Society of America* (2003).

Dilek, Yildirim, and Harald Furnes. "Ophiolite genesis and global tectonics: Geochemical and tectonic fingerprinting of ancient oceanic lithosphere." *Bulletin* 123, no. 3-4 (2011): 387-411.

Dodd, MS, et al. (2017) Evidence for early life in Earth's oldest hydrothermal vent precipitates. Nature, 543 (7643). pp. 60-64.

Eaton, Jerry P. (1963). Crustal structure from San Francisco, California, to Eureka, Nevada, from seismic-refraction measurements, Journal of Geophysical Research.

Elder, Will. (2013). Bedrock geology of the San Francisco Bay Area: A local sediment source for bay and coastal systems. Marine Geology. 345. 18-30.

Ellsworth, W.L., Beroza, G.C., Julian, B.R., Klein, F., Michael, A.J., Oppenheimer, D.H., Prejean, S.G., Richards-Dinger, K., Ross, S.L., Schaff, D.P., and Waldhauser, F., 2000, Seismicity of the San Andreas fault system in central California: Application of the double-difference location algorithm on a regional scale: Eos, Transactions, American Geophysical Union, v. 81, p. F919.

Faust, G.T. and Fahey, J.J. (1962). The Serpentine-Group Minerals Studies Of The Natural Phases In The System Mgo-Sio2-H2o And The Systems Containing The Congeners Of Magnesium, Geological Survey Professional Paper 384-A.

Ferriz, H. and Anderson, R. (2001). Engineering Geology Practice In Northern California, State of California Resources Agency, Dept. of Conservation.

Frost, B. Ronald, and James S. Beard. "On Silica Activity and Serpentinization." *Journal of Petrology* 48, no. 7 (2007): 1351-1368.

Früh-Green, G. L., et al, "30,000 Years of Hydrothermal Activity at the Lost City Vent Field." *Science* 301, no. 5632 (2003): 495-498.

Furnes, H. and Dilek, Y. Archean versus Phanerozoic oceanic crust formation and tectonics: Ophiolites through time, *Geosystems and Geoenvironment,* Volume 1, Issue 1, (February 2022).

Galloway, J.P. et al, (2001). Consumer Uses of Industrial Minerals in the San Francisco Bay Area—Houses to Interstates, Geology and Natural History of the San Francisco Bay Area, 2001 Fall Field Conference.

Gardner, R.A. et al, U.S. Dept. of Agriculture, Soil Survey of the Santa Clara Area California (June 1958).

Geological Gems of California State Parks, Special Report 230 – Fuller, M., Brown, S., Wills, C. and Short, W., editors, 2015, Geological Gems of California, California Geological Survey Special Report 230 under Interagency Agreement C01718011 with California State Parks.

Giaramita, Mario, Glenn J. MacPherson, and Stephen Paul Phipps. "Petrologically diverse basalts from a fossil oceanic forearc in California: The Llanada and Black Mountain remnants of the Coast Range ophiolite." *Geological Society of America Bulletin* 110, no. 5 (1998): 553-571.

Gifford, Jennifer & Malone, Shawn & Mueller, Paul. (2020). The Medicine Hat Block and the Early Paleoproterozoic Assembly of Western Laurentia. *Geosciences*. 10. 271.

Gjøvik, Ashley M. "The Lost Hydrothermal Crystals of Santa Clara County: A Structural and Archival Study of Six Rare Minerals from a Single Geologic Province." *The Journal of Decolonized Ecology and Evolution* 1, no. 1 (June 9, 2025). https://doi.org/10.5281/zenodo.15626525.

Gjøvik, Ashley, M. "Compiled Chemistry: Structural Inheritance and the Deep Origins of Biological Function". *The Journal of Decolonized Ecology and Evolution* 1, no. 1 (June 23, 2025). https://doi.org/10.5281/zenodo.15717955.

Gjøvik, Ashley M. "Thermal, UV, and Impact Constraints on Pre-cambrian Terrestrial Evolution: The Impossibility of Complex Multicellular Life on Land Prior to 600 MYA". *The Journal of Decolonized Ecology and Evolution* 1, no. 1 (June 25, 2025). https://doi.org/10.5281/zenodo.15739885.

Godfrey, Nicola J., and Klemperer, Simon. L. Ophiolitic basement to a forearc basin and implications for continental growth: The Coast Range/Great Valley ophiolite, California. *Tectonics*, Vol. 17, No. 4, Pages 558-570, August 1998.

Godfrey, Nicola J., and Yildirim Dilek. "Mesozoic assimilation of oceanic crust and island arc into the North American continental margin in California and Nevada: Insights from geophysical data." (2000).

Graymer, Russell & Givler, Robert & Baldwin, J. & Lettis, William & Johnson, Samuel & Greene, H. & Dartnell, Peter. (2018). Geology of San Francisco.

Graymer, R. & Stanley, Richard & Ponce, David & Jachens, R. & Simpson, R. & Wentworth, C. (2015). Structural superposition in fault systems bounding Santa Clara Valley, California. Geosphere. 11. 63-75.

Grossinger, R. et al, Prepared for the Santa Clara Valley Water District and The Nature Conservancy, South Santa Clara Valley Historical Ecology Study, May 2008.

Gurdak, J. and Cook, G. 2021 Groundwater Management Plan for the Santa Clara and Llagas Subbasins, Santa Clara Valley Water District, Nov. 2021.

Guillaume Bonnet, F E Apen, Mathieu Soret, Julie Noel, Benoît Caron, et al. Seamount Subduction Dynamics and Long-Term Evolution of the Franciscan Active Margin. Tectonics, 2024, 43 (3), pp.e2023TC008084.

Gustafsson, Jacob & Drake, Henrik. (2025). Biosignatures of an ancient bedrock- and impact structure-hosted deep biosphere: current knowledge and future perspectives. *Discover Geoscience.* 3.

Hanson, Randall. (2015). Hydrologic framework of the Santa Clara Valley, California. *Geosphere*.

Hausel, W. D. Pacific Coast Diamonds – An Unconventional Source Terrane. Mineral Report MR94-8, *Wyoming State Geological Survey*.

Healy, J.H. (1963). Crustal structure along the coast of California from seismic-refraction measurements, *Journal of*

Geophysical Research, v. 68, no. 20.

Heron, Philip J., J. Brendan Murphy, R. Damian Nance, and Russell N. Pysklywec. “Pannotia's mantle signature: the quest for supercontinent identification.” *Special Publications* 503 (2020): 41 – 61.

Himson, Stephen, et al. (2023). The San Francisco Estuary, USA as a reference section for an Anthropocene series. *The Anthropocene Review*. 10.

Hodgson, Susan F. (1987). Onshore Oil & Gass Seeps in California, California Department of Conservation, Division of Oil & Gas, Publication No. TR26, 1980-1987.

Holm NG, Oze C, Mousis O, Waite JH, Guilbert-Lepoutre A. Serpentinization and the Formation of H2 and CH4 on Celestial Bodies (Planets, Moons, Comets). *Astrobiology*. 2015;15(7):587-600.

Holmes, L.C. and Nelson, J.W. and Lapham, M.H. (1917). U.S. Department of Agriculture, Reconnaissance Soil Survey of the San Francisco Bay Region California.

Horodyskyj, Ulyana & Lee, Cin-Ty & Ducea, Mihai. (2007). Similarities between Archean high MgO eclogites and Phanerozoic arc-eclogite cumulates and the role of arcs in Archean continent formation. Earth and Planetary Science Letters. 256. 510-520.

Hu, D. (2014) The Model of Oceanic Crust Expansion. *International Journal of Geosciences*, 5, 1594-1601.

Huot, F. and Muary, R.C. The Round Mountain serpentinite mélange, northern Coast Ranges of California: An association of backarc and arc-related tectonic units, *GSA Bulletin* (2002) 114 (1): 109–123.

Jachens, R.C., and Moring, B.C., 1990, Maps of the thickness of Cenozoic deposits and the isostatic residual gravity over basement for Nevada: U.S. Geological Survey Open-File Report 90-404, 15 p., 2 sheets, scale 1:1,000,000.

Jachens, R.C., Wentworth, C.M., and McLaughlin, R.J., 1998, Pre-San Andreas location of the Gualala block inferred from magnetic and gravity anomalies, *in* Elder, W.P., (ed.), Geology and tectonics of the Gualala block, northern California: The Pacific Section, SEPM, Society of Sedimentary Geology, Book 84, p. 27-64.

Jachens, R., Simpson, R., Graymer, R., Wentworth, C., and Brocher, T., 2006, Three-dimensional geologic map of northern and central California: A basic model for supporting earthquake simulations and other predictive modeling [abs.]: Seismological Research Letters, v. 77, p. 271.

Jachens, R.C. et al. (2017). The Evergreen basin and the role of the Silver Creek fault in the San Andreas fault system, San Francisco Bay region, Californ. *Geosphere*. 13.

Jachens, R.C. et al, (2001), 3D Geologic Maps and Visualization: A New Approach to the Geology of the Santa Clara (Silicon) Valley, California, Digital Mapping Techniques '01, Workshop Proceedings U.S. Geological Survey Open-File Report 01-223.

Ingersoll, Raymond V. "Models for origin and emplacement of Jurassic ophiolites of northern California." (2000).

Keppie, J.D. et al, 2004, Mexican data indicating ca.700 Ma breakup of Rodinia and ca.550 Ma separation of Avalonia: analogous to events in western Laurentia, American Geophysical Union, Spring Meeting 2004, abstract id. GP23A-01.

Kerrich, R. & Jia, Yongheng & Wyman, Derek. (2001). Transition From Archean Plume-Arc Orogens to Phanerozoic Style Convergent Margin Orogens and Changing Mantle Lithosphere. AGU Fall Meeting Abstracts.

Kirscher, Uwe & et al. (2020). Paleomagnetic Constraints on the Duration of The Australia-Laurentia Connection in the Core of the Nuna Supercontinent. Geology. 49.

Klein, Frieder, and Wolfgang Bach. "Serpentinization." In *Encyclopedia of Marine Geosciences*, edited by Jan Harff, Martin Meschede, Sven Petersen, and Jörn Thiede. Dordrecht: Springer, 2009.

Klein, Frieder, et al. "Geological Evidence for High H_2 Production from Komatiites in the Archaean." *Nature Geoscience* 17 (2023): 1-7.

Karlstrom, K.E., Ahall, K.-I., Harlan, S.S., William, M.L., McLelland, J., and Geissman, J.W., 2001, Long-lived (1.8–0.8 Ga) convergent orogen in southern Laurentia, its extensions to Australia and Baltica, and implications for refining Rodinia: *Precambrian Research*, v. 111 p. 5-30.

Kassab, B. and Cook, G. 2016. Groundwater Management Plan Santa Clara and Llagas Subbasins. Santa Clara Valley Water District, (November 2016).

Keefer, David. (1998). The Loma Prieta, California, Earthquake of October 17,1989 Landslides, U.S. Geological Survey Professional Paper 1551-C.

Kimura, G. et al, Well-preserved Underplating Structure of the Jadeitized Franciscan Complex, Pacheco Pass, California, *Geology*; January 1996; v. 24; no. 1; p. 75–78.

Kious, J. W. and Tilling, R. I., 1996, This Dynamic Earth: The Story of Plate Tectonics: USGS General Interest Publication.

Kirby, S.H., Wang, K. & Brocher, T.M. A large mantle water source for the northern San Andreas fault system: a ghost of subduction past. *Earth Planet Sp* **66**, 67 (2014).

Kirby. S. H. et al., "Evidence for multiple stages of serpentinization from the mantle through the crust in the Redwood City Serpentinite mélange along the San Andreas Fault in California," *Lithos* 336-337 (2019): 16-36.

Kirscher, U., Mitchell, R.N., Liu, Y., Pisarevsky, S.A., Giddings, J., Li, Z.X., 2022. Paleomagnetic evidence for a Paleoproterozoic rotational assembly of the North Australian Craton in the leadup to supercontinent formation. Geophys. Res. Lett. 49(22).

Klein, Frieder, et al. "Carbonation of Serpentinite in Creeping Faults of California." *Geophysical Research Letters* 49, no. 18 (2022): e2022GL099185.

Krohe, Alexander. (2017). GR Focus Review Article: The Franciscan Complex (California, USA) – the model case for return-flow in a subduction channel put to the test. Gondwana Research. 45. 282-307.

Kusky TM, Celâl Şengör AM. Comparative orotomy of the Archean Superior and Phanerozoic Altaid orogenic systems. *Natl Sci Rev.* 2022;10(2):nwac235.

Kusky, Timothy & Li, Jianghai. (2010). Origin and Emplacement of Archean Ophiolites of the Central Orogenic Belt, North China Craton. *Journal of Earth Science.* 21. 744-781.

Kusky, Timothy & Windley, Brian & Polat, Ali & Wang, Lu & Ning, Wenbin. (2021). Archean dome-and-basin style structures form during growth and death of intraoceanic and continental margin arcs in accretionary orogens. *Earth-Science Reviews.* 220.

Langenheim, V. E. et al, A summary of the late Cenozoic stratigraphic and tectonic history of the Santa Clara Valley, California, *Geosphere*; February 2015; v. 11; no. 1; p. 50–62.

Langenheim, V.E., Schmidt, K.M., and Jachens, R.C., 1997, Coseismic deformation during the 1989 Loma Prieta earthquake and range-front thrusting along the southwestern margin of the Santa Clara Valley, California: *Geology*, v. 25, p. 1091–1094.

Langenheim, Victoria & Jachens, R. & Wentworth, C. & Mclaughlin, R. (2013). Previously unrecognized regional structure of the Coastal Belt of the Franciscan Complex, Northern California, revealed by magnetic data. *Geosphere*. 9. 1514-1529.

Leidy, R.A., G.S. Becker, B.N. Harvey. 2005. Historical distribution and current status of steelhead/rainbow trout (Oncorhynchus mykiss) in streams of the San Francisco Estuary, California. Center for Ecosystem Management and Restoration, Oakland, CA

Leong, James Andrew M., Tucker Ely, and Everett L. Shock. "Decreasing Extents of Archean Serpentinization Contributed to the Rise of an Oxidized Atmosphere." *Nature Communications* 12, no. 1 (2021): 7341.

Levy, M. and Christie-Blick, N., Nicholas, 1991. Tectonic subsidence of the early Paleozoic passive continental margin in eastern California and southern Nevada. Geological Society of America Bulletin, v. 103, p. 1590–1606.

Li, Z-X. et al, Assembly, configuration, and break-up history of Rodinia: A synthesis, *Precambrian Research,* 160, 1-2, 2008.

Little, Crispin & Herrington, R. & Haymon, Rachel & Danelian, Taniel. (1999). Early Jurassic hydrothermal vent community from the Franciscan Complex, San Rafael Mountains, California. Geology. 27.

Liu, Fei, Dongyang Lian, Weiwei Wu, and Jingsui Yang. 2021. "Diamond and Other Exotic Mineral-Bearing Ophiolites on the Globe: A Key to Understand the Discovery of New Minerals and Formation of Ophiolitic Podiform Chromitite" *Crystals* 11, no. 11: 1362.

Lever, J. P., Sundell, K. E., Pearson, D. M., & Brennan, D. T. (2024). Recently identified Mesoproterozoic strata in south-central Idaho document late-stage rifting of the Nuna supercontinent in western Laurentia. Geochemistry, Geophysics, Geosystems, 25, e2024GC011503

Levy, M. & Christie-Blick, N. (1991). Tectonic subsidence of the early Paleozoic passive continental margin in eastern California and southern Nevada. Geological Society of America Bulletin. 103. 1590-1606.

Lockwood, J.P. "Sedimentary and gravity-slide emplacement of serpentinite." *Geological Society of America Bulletin* 82, no. 4 (1971): 919-936.

Macdonald, F. A., et al., 2013. The Laurentian record of Neoproterozoic glaciation, tectonism, and eukaryotic evolution in Death Valley, California. Geological Society of America Bulletin, v. 125, p. 1203–1223.

Malpas, J. (1992). Serpentine and the geology of serpentinized rocks. In: Roberts, B.A., Proctor, J. (eds) The Ecology of Areas with Serpentinized Rocks. Geobotany, vol 17. Springer, Dordrecht.

McLaughlin, R.J., Clark, J.C., and Brabb, E.E., 1988, Geologic map and structure sections of the Loma Prieta 7 Vi′ Quadrangle, Santa Clara and Santa Cruz counties, California: U.S. Geological Survey Open- File Map 88-752, 32 p., scale 1:24,000, 2 sheets.

McLaughlin, R.J., Clark, J.C., Brabb, E.E., and Helley, E.J., 1991, Geologic map and structure sections of the Los Gatos *l- ll2* Quadrangle, Santa Clara and Santa Cruz counties, California: U.S. Geological Survey Open-File Map 91-593, 48 p., scale 1:24,000, report and three plates.

McLaughlin, R.J., et al., 2000, Neogene contraction between the San Andreas fault and the Santa Clara Valley, San Francisco Bay Region, California, *in* Ernst, W.G., and Coleman, R.G., (eds.), Tectonic studies of Asia and the Pacific rim: Geological Society of America, International Book Series, v. 3, p. 265-294.

McLaughlin, R.J. et al, (2001). Geologic Maps and Structure Sections of the Southwestern Santa Clara Valley and Southern Santa Cruz Mountains, Santa Clara and Santa Cruz Counties, California, U.S. Geological Survey Miscellaneous Field Studies MF–2373.

McLaughlin, Robert J., and Benjamin L. Melosh. "Integrated geologic and geophysical modeling across the Bartlett Springs fault zone, northern California (USA): Implications for fault creep and regional structure." *Geosphere* 20, no. 1 (2024): 129-158.

Medley, Edmund and Zekkos, Dimitrios. (2011). *Geopractitioner approaches to working with antisocial mélanges.* Mélanges: Processes of Formation and Societal Signifi cance: Geological Society of America Special Paper 480, p. 261–277.

Medley, Edmund. (2008). Engineering the geological chaos of franciscan and other bimrocks. 42nd U.S. Rock Mechanics - 2nd U.S.-Canada Rock Mechanics Symposium.

Meert, J.G. "A synopsis of events related to the assembly of eastern Gondwana," *Tectonophysics* 362, no. 1-4 (2003): 1-40.

Merdith, Andrew S., Sascha Brune, Anne Glerum, and Douwe J.J. van Hinsbergen. "Global Hydrogen Production During High-Pressure Serpentinization of Subducting Slabs." *Geochemistry, Geophysics, Geosystems* 24, no. 3 (2023): e2023GC010947.

Metropolitan Transportation Commission & Association of Bay Area Governments, Plan Bay Area 2050, Draft EIR: 3.5 Biological Resources (June 2021).

Metropolitan Transportation Commission & Association of Bay Area Governments, Plan Bay Area 2050, Draft EIR: 3.8 Geology, Seismicity, and Mineral Resources (June 2021).

Metropolitan Transportation Commission & Association of Bay Area Governments, Plan Bay Area 2050, Draft EIR: 3.10 Hydrology and Water Quality (June 2021).

Mohan, Swetha, et al. "Serpentine alteration as source of high dissolved silicon and elevated δ30Si values to the marine Si cycle." *Nature Communications* 11, no. 1 (2020): 5854.

Monnin, Christophe, et al. (2021). The Chemistry of Hyperalkaline Springs in Serpentinizing Environments: 1. The Composition of Free Gases in New Caledonia Compared to Other Springs Worldwide. *Journal of Geophysical Research: Biogeosciences.* 126.

Moore, D. E., Metamorphic History of a High-Grade Blueschist Exotic Block from the Franciscan Complex, California, *Journal of Petrology*, Volume 25, Issue 1, February 1984, Pages 126–150.

Moores, E.M. and Jackson, E.D. (1974). Ophiolites and oceanic crust. *Nature*.

Morrill, Penny L., et al. "Geochemistry and geobiology of a present-day serpentinization site in California: The Cedars." *Geochimica et Cosmochimica Acta* 109 (2013): 222-240.

Mulcahy, S. R. Early Onset of Franciscan Subduction, *Tectonics*. Volume 37, Issue 5, May 2018, Pages 1194-1209.

Munro-Fraser, J. P. (1881), "History of Santa Clara County, California: including its geography, geology, topography, climatography and description, together with a record of the Mexican grants; its mines and natural springs…" Alley, Bowen & Co. Publishers, San Francisco.

Murphy, J. B. "Pannotia: in defence of its existence and geodynamic significance." *Special Publications* 503 (2020): 13 - 39.

Nance RD. The supercontinent cycle and Earth's long-term climate. *Ann N Y Acad Sci*. 2022;1515(1):33-49.

Nelson, L. L., et al., 2020. Geochronological constraints on Neoproterozoic rifting and onset of the Marinoan glaciation from the Kingston Peak Formation in Death Valley, California (USA). Geology, v. 48, p. 1083–1087.

Nordsvan, A. et al. (2018). Laurentian crust in northeast Australia: Implications for the assembly of the supercontinent Nuna. Geology. 46.

Nordsvan, A., et al. (2022). Post-collisional magmatism in NE Australia during Mesoproterozoic supercontinent Nuna: Insights from new zircon U Pb and Lu Hf data. Lithos. 428-429.

Ogawa, Y., Mori, R., Tsunogae, T., Dilek, Y., & Harris, R. (2014). New interpretation of the Franciscan mélange at San Simeon coast, California: tectonic intrusion into an accretionary prism. *International Geology Review*, *57*(5–8), 824–842.

Oguadinma, Vivian & Kelechi, Ibekwe & Lanisa, Ademola & Victory, Okoro & Victor, Ahaji & Chukwudalu, Ezeonyema & Chinazaepkere, Arukwe. (2023). Submarine canyon: A brief review.

O'Neil, James R., and Hugh P. Taylor. "Stable-isotope studies of rocks and secondary minerals in a vapor-dominated hydrothermal system at The Geysers, Sonoma County, California." *Geochimica et Cosmochimica Acta* 53, no. 8 (1989): 2025-2039.

Oze, Christopher. (2004). Chromium geochemistry in serpentinized ultramafic rocks and serpentine soils from the Franciscan complex of California. American Journal of Science - AMER J SCI. 304. 67-101.

Ponti, D.J., and Wells, R.E., in press, Off fault ground ruptures in the Santa Cruz Mountains, California: Ridge-top spreading vs. tectonic extension during the 1989 Loma Prieta earthquake: Seismological Society of America Bulletin.

Parsons, Tom. (1998). Seismic-reflection evidence that the Hayward fault extends into the lower crust of the San Francisco Bay area, California. *Bulletin of the Seismological Society of America*. 88. 1212-1223.

Pehrsson, S. et al. “Metallogeny and its link to orogenic style during the Nuna supercontinent cycle.” *Special Publications* 424 (2015): 83 - 94.

Pourteau, A., et alJ., 2018. 1.6 Ga crustal thickening along the final Nuna suture. Geology 46(11), 959-962.

Powell, C. L. Age, Stratigraphy, and Correlations of the Late Neogene Purisima Formation, Central California Coast Ranges. U.S. Geological Survey Professional Paper 1740.

Prodehl, Claus. (1979). Crustal Structure of the Western United States, Geological Survey Professional Paper 1034.

Rasmussen KL, Stamps BW, Vanzin GF, Ulrich SM, Spear JR. Spatial and temporal dynamics at an actively silicifying hydrothermal system. *Front Microbiol*. 2023;14:1172798.

Raymond, Loren A. 2019. "Origin of Mélanges of the Franciscan Complex, Diablo Range and Northern California: An Analysis and Review" *Geosciences* 9, no. 8: 338.

Raymond, Loren & Bero, David. (2023). Paleogeographic reconstruction of regional accretionary complex architecture, Franciscan Complex, northwestern San Francisco Bay Area, California, USA. Geosphere. 19.

Raymond, L. A. (2024). Deconstruction of the Franciscan Complex Central Terrane Mélange and re-evaluation of Franciscan mélanges and architecture of the northwestern San Francisco Bay Area, California, USA, *Journal of the Geological Society*.

Rejeck, Peter. July 25 2008. 'One in a million': Granite boulder links Antarctica to western United States as part of ancient supercontinent, NSF U.S. Antarctica Program.

Roberts, Carter W., Jachens, Robert C., 2003, Shaded Relief Aeromagnetic Map of the Santa Clara Valley and Vicinity, California: *U. S. Geological Survey,* Open-File Report 03-360.

Roberts, C.W., Jachens, R.C., Oliver, H.W., (1990). Isostatic Residual Gravity Map of California and Offshore Southern California, California Geological Data Map Series No. 7.

Robbins, S.L., Oliver, H.W., and Holden, K.D. (1976). Bouguer Gravity Map of California: San Jose Sheet, California Division of Mines & Geology.

Robbins, S.L. and Roberts, C.W., (1982), Boguer gravity map of the southern San Francisco Bay region, California, U.S. Geological Survey, Open-File Report 78-733.

Robinson, Paul & Malpas, John & Dilek, Yildirim & Zhou, Mei-Fu. (2008). The significance of sheeted dike complexes in ophiolites. *GSA Today*. 18.

Rutte, D. et al. (2020). An Exhumation Pulse From the Nascent Franciscan Subduction Zone (California, USA). *Tectonics*. 39.

Rytuba, J.J. et al, Genesis of Silica-Carbonate Type Mercury Ore Deposits in Coast Range California from Mantle Derived Fluids, American Geophysical Union, Fall Meeting 2016, abstract #DI23B-2612 (Dec. 2016).

Rytuba, James. J. Mercury from mineral deposits and potential environmental impact, *Environmental Geology* (2003) 43:326–338.

Sanders, Ian L. (2012). The Mineral Springs of Santa Clara County, San Martin, California.

Santa Clara City, Planning Division, CEQA Archive Santa Clara Square Residential/Mixed-Use Project, 2015-2017, https://www.santaclaraca.gov/Home/Components/BusinessDirectory/BusinessDirectory/121/3650

Santa Clara County, Department of Planning and Development, County Of Santa Clara Historic Context Statement, December 2004 (Revised February 2012).

Santa Clara Valley Urban Runoff Pollution Prevention Program (SCVURPPP). 2018. "San Tomas Aquino Watershed." April 26, 2018.

San Francisco Estuary Institute, "West Valley Watershed Assessment 2018: Baseline Ecological Condition Assessment of Southwest San Francisco Bay Creeks in Santa Clara County; Calabazas, San Tomas Aquino, Saratoga, Sunnyvale East and West," SFEI Contribution No. 944 (2019).

SCBWMI. 2001. Watershed management plan, Volume One unabridged, watershed characteristics report, Chapter 7: natural setting (amended December 2003). Santa Clara Basin Watershed Management Initiative.

SCBWMI. 2003. Fact Sheet No. 1: A Publication for the

Santa Clara Basin Watershed Management Initiative, The Santa Clara Basin, (August 2003).

Schmidt, D. A., and R. Bürgmann (2003), Time-dependent land uplift and subsidence in the Santa Clara valley, California, from a large interferometric synthetic aperture radar data set, *J. Geophys. Res.*, 108, 2416.

Schmidt, K.M. et al, (2014). Deformation from the 1989 Loma Prieta earthquake near the southwest margin of the Santa Clara Valley, California, *Geosphere*; December 2014; v. 10; no. 6; p. 1177–1202.

Schulte, Mitchell, Daniel Blake, Thomas Hoehler, and Timothy McCollom. "Serpentine and Serpentinization: A Link Between Planet Formation and Life." *Geology* 38, no. 10 (2010): 959-960.

Scott, E.S. (2016). In My Backyard: Creationism in California, National Center for Science Education.

Segall, P. and Lisowski, M, 1990, Surface displacements in the 1906 San Francisco and 1989 Loma Prieta earthquakes: Science, v. 250, p. 1241-1244.

Shervais, J. W., 2001. Birth, death, and resurrection: The life cycle of suprasubduction zone ophiolites, *Geochem. Geophys. Geosyst.,* vol. 2.

Shervais, J. et al, (2004) Multi-Stage Origin of the Coast Range Ophiolite, California: Implications for the Life Cycle of Supra-Subduction Zone Ophiolites, *International Geology Review*, Vol. 46, 2004, p. 289–315.

Shervais, J. et al, (2005). Radioisotopic and biostratigraphic age relations in the Coast Range Ophiolite, northern California: Implications for the tectonic evolution of the Western Cordillera. *GSA Bulletin*; May/June 2005; v. 117; no. 5/6; p. 633–653

Shervais, John. (2006). Plumes, Plateaux, and the Significance of Subduction-related Accretionary Complexes in Early Earth Processes. Special Paper of the Geological Society of America. 405.

Shervais, John W. "Tonalites, trondhjemites, and diorites of the Elder Creek ophiolite, California: low-pressure slab melting and reaction with the mantle wedge." (2008).

Shervais, John W., et al, (2011), Serpentinite matrix mélange: Implications of mixed provenance for mélange formation, *Mélanges: Processes of Formation and Societal Significance,* Geological Society of America.

Shervais, John W. and Choi, S. H. (2011), Subduction initiation along transform faults: The proto-Franciscan subduction zone. Lithosphere; v. 4; no. 6; p. 484–496.

Shervais, John W. and Choi, S. H. (2012), Subduction initiation along transform faults: The proto-Franciscan subduction zone, Lithosphere (2012) 4 (6): 484–496.

Snortum, Eric & Day, James. (2020). Forearc origin for Coast Range Ophiolites inferred from osmium isotopes and highly siderophile elements. Chemical Geology. 550. 119723.

Snow, C. A., Wakabayashi, J., Ernst, W. G., & Wooden, J. L. (2010). Detrital zircon evidence for progressive underthrusting in Franciscan metagraywackes, west-central California. *Bulletin*, *122*(1-2), 282-291.

Sorg, D.H., and McLaughlin, R.J, 1975, Geologic Map of the Sargent-Berrocal fault zone between Los Gatos and Los Altos Hills, Santa Clara County, California: U.S. Geological Survey Miscellaneous Field Investigations Map MF-643, scale 1:24,000.

Stanley, P. G., et al. (2002). Subsurface and petroleum geology of the southwestern Santa Clara Valley ("Silicon Valley"), California. U.S. Geological Survey, USGS Numbered Series 1663.

Stanley, R.G., Jachens, R.C., Kvenvolden, K.A., Hostettler, F.D., Magoon, L.B., and Lillis, P.G., 1996, Evidence for an oil-bearing sedimentary basin of probable Miocene age beneath "Silicon Valley", California: annual Meeting Abstracts-American Association of Petroleum Geologists and Society of Economic Paleontologists and Mineralogists, v. 5, p.133-134.

Stern, Bob. (2005). Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that the modern episode of subduction tectonics began in Neoproterozoic time. *Geology*. 33.

Stewart, J. H. Reconstructing Rodinia by Fitting Neoproterozoic Continental Margins, Open-File Report 2009-1191, U.S. Geological Survey.

Sossi, Paolo A., Stephen M. Eggins, Robert W. Nesbitt, Oliver Nebel, Janet M. Hergt, Ian H. Campbell, Hugh St.C. O'Neill, Martin Van Kranendonk, and D. Rhodri Davies. "Petrogenesis and Geochemistry of Archean Komatiites." *Journal of Petrology* 57, no. 1 (2016): 147-184.

Stoffer, Philip. Rocks and Geology in the San Francisco Bay Region, U.S. Geological Survey, Bulletin 2195 (2002).

Swanson-Hysell, Nicholas, et al. (2021). The Paleogeography of Laurentia in Its Early Years: New Constraints From the Paleoproterozoic East-Central Minnesota Batholith. Tectonics. 40.

Swanson-Hysell, Nicholas. (2021). The Precambrian paleogeography of Laurentia. 10.1016/B978-0-12-818533-9.00009-6.

Taliaferro, Nicholas L.; Franciscan-Knoxville Problem. *AAPG Bulletin* 1943; 27 (2): 109–219.

Teixeira, João & Misi, Aroldo & Silva, Maria. (2007). Supercontinent evolution and the Proterozoic metallogeny of South America. Gondwana Research - GONDWANA RES. 11. 346-361.

Tsujimori, T., Matsumoto, K., Wakabayashi, J. *et al.* Franciscan eclogite revisited: Reevaluation of the *P*–*T* evolution of tectonic blocks from Tiburon Peninsula, California, U.S.A. *Mineralogy and Petrology* **88**, 243–267 (2006).

Wagner, D. L. et al, 2002, Tar Creek study, Sargent oil field, Santa Clara County, California, California Dept. of Conservation, Division of Oil, Gas, and Geothermal Resources.

Wakabayashi, John. (1992). Nappes, Tectonics of Oblique Plate Convergence, and Metamorphic Evolution Related to 140 Million Years of Continuous Subduction, Franciscan Complex, California. The Journal of Geology, Vol. 100, No. 1 (Jan., 1992), pp. 19-40.

Wakabayashi, John. (2008). Franciscan complex, California: Problems in recognition of melanges, and the gap between research knowledge and professional practice. 42nd U.S. Rock Mechanics - 2nd U.S.-Canada Rock Mechanics Symposium.

Wakabayashi, John. (2011). Mélanges of the Franciscan Complex, California: Diverse structural setting, evidence for sedimentary mixing, and their connection to subduction processes. Special Paper of the Geological Society of America. 480. 117-141.

Wakabayashi, J. (2015). Anatomy of a subduction complex: architecture of the Franciscan Complex, California, at multiple length and time scales. *International Geology Review*, *57*(5–8), 669–746.

Wakabayashi, John & Rowe, Christie. (2015). Whither the megathrust? Localization of large-scale subduction slip along the contact of a mélange. International Geology Review. 57. 854-870.

Wakabayashi, J. (2017). Structural context and variation of ocean plate stratigraphy, Franciscan Complex, California: insight into mélange origins and subduction-accretion processes. *Prog Earth Planet Sci* **4**, 18.

Wakabayashi, John. (2021). Field and petrographic reconnaissance of Franciscan complex rocks of Mount Diablo, California: Imbricated ocean floor stratigraphy with a roof exhumation fault system. *Regional Geology of Mount Diablo, California: Its Tectonic Evolution on the North America Plate Boundary.* Geological Society of America.

Wentworth, C.M. et al. (2005). Quaternary Deposits of the Santa Clara Valley, California. San Jose GSA, AAPG Joint Meeting.

Wesling, J.R. and Helley, E.J. (1989), Quaternary geologic map of the San Jose West Quadrangle, Santa Clara County, California, Open-File Report 89-672, U.S. Dept. of the Interior, Geological Survey.

Whitmeyer, Steven & Karlstrom, Karl. (2007). Tectonic model for the Proterozoic growth of North America. Geosphere. 3. 220.

Ueda, H., et al. "Chemical Nature of Hydrothermal Fluids Generated by Serpentinization and Carbonation of Komatiite: Implications for H_2-Rich Hydrothermal System and Ocean Chemistry in the Early Earth." *Geochemistry, Geophysics, Geosystems* 22, no. 11 (2021).

U.S. Department of Agriculture, Natural Resources Conservation Service. 2015. Supplement to the Soil Survey of the Santa Clara Area, California, Western Part.

U.S. Environmental Protection Agency (Feb. 2005), Proceedings and Summary Report: Workshop on Assessing and Managing Mercury from Historic and Current Mining Activities, Nov. 28 – 30, 2000, San Francisco California, EPA/625/R-04/102.

U.S. Geological Survey. "Shaded relief aeromagnetic map of the Santa Clara Valley and vicinity, California." Open-File Report 03-360. Reston, VA: USGS, 2003.

U.S. Geological Survey. "Geology of the Southernmost Part of Santa Clara County, California: A Digital Database." Open-File Report 97-710. Reston, VA: USGS, 1997.

U.S. Geological Survey. "Subsurface geometry of the San Andreas-Calaveras fault junction: Influence of serpentinite and the Coast Range Ophiolite." Professional Paper 1737. Reston, VA: USGS, 2004.

U.S. Geological Survey. "Crystal Knob volcanic pipe peridotite xenoliths and mantle lithosphere structure." Bulletin 2206. Reston, VA: USGS, 2006.

U.S. Geological Survey. "Modeling and validation of a 3D velocity structure for the Santa Clara Valley, California, for seismic-wave simulations." Professional Paper 1758. Reston, VA: USGS, 2007.

Ukar, Estibalitz, Cloos, Mark, and Vasconcelos, Paulo. First ^{40}Ar-^{39}Ar Ages from Low-T Mafic Blueschist Blocks in a Franciscan Mélange near San Simeon: Implications for Initiation of Subduction , *The Journal of Geology* (2012) 120 (5): 543–556.

Ukar, Estibalitz. (2012). Tectonic significance of low-temperature blueschist blocks in the Franciscan mélange at San Simeon, California. Tectonophysics. s 568–569. 154–169.

Ukar, Estibalitz & Cloos, Mark. (2019). Cataclastic deformation and metasomatism in the subduction zone of mafic blocks-in-mélange, San Simeon, California. Lithos. 346-347.

Uno, Masaoki & Kirby, Stephen. (2019). Evidence for multiple stages of serpentinization from the mantle through the crust in the Redwood City Serpentinite mélange along the San Andreas Fault in California. Lithos. 336.

Verbaas, J.; et al. (2018). "Rifting of Western Laurentia at 1.38 Ga: The Hart River Sills of Yukon, Canada". *Lithos, 316-317*, 243-260.

Volante, Silvia & Collins, William & Pourteau, Amaury & Li, Zheng-Xiang & Li, Jiangyu & Nordsvan, Adam. (2020). Structural Evolution of a 1.6 Ga Orogeny Related to the Final Assembly of the Supercontinent Nuna: Coupling of Episodic and Progressive Deformation. *Tectonics*. 39.

Volante S, Kirscher U. (2024). Metamorphic turnover at 2 Ga related to two-stage assembly of Columbia. *Sci Rep*. 2024;14(1):6483.

Walkinshaw, Mike, A.T. O'Geen, D.E. Beaudette. "Soil Properties." *California Soil Resource Lab*, 1 Oct. 2023, casoilresource.lawr.ucdavis.edu/soil-properties/.

Wagner, David. L. et al, Tar Creek Study, Sargent Oil Field Santa Clara County, California, Cal. Dept. of Conservation, Open-file Report No. 5, 2002.

Wang, Y., et al. (2019). The exhumation of subducted oceanic-derived eclogites: Insights from phase equilibrium and thermomechanical modeling. *Tectonics*, 38, 1764–1797.

Waterton, Pedro & Arndt, Nicolas. (2023). Komatiites: their geochemistry and origins. The Archaean Earth: Tempos and Events (2nd Edition of The Precambrian Earth). Elsevier.

Wenner, David B., and Hugh P. Taylor. "Oxygen, hydrogen and carbon isotope studies of the Franciscan formation, Coast Ranges, California." *Geochimica et Cosmochimica Acta* 40, no. 7 (1976): 721-734.

Wentworth, C.M., Blake, M.C., Jr., McLaughlin, R.J., and Graymer, R.W., 1998, Preliminary geologic map of the San Jose 30 X 60-minute quadrangle, California: a digital database: U.S. Geological Survey Open-File Report 98-795.

Wentworth, C.M., Williams, R.A., Jachens, R.C., Graymer, R.W., and Stephenson, W.J., 2010, The Quaternary Silver Creek Fault beneath the Santa Clara Valley, California: U.S. Geological Survey Open-File Report, 2010-1010, 50 p.

Whitmeyer, Steven J. and Karl E. Karlstrom. "Tectonic model for the Proterozoic growth of North America." *Geosphere* 3 (2007): 220-259.

Xia, Yan & Xu, Xisheng. (2019). A Fragment of Columbia Supercontinent: Insight for Cathaysia Block Basement From Tectono-Magmatic Evolution and Mantle Heterogeneity. Geophysical Research Letters. 46.

Yancey, T., Lee, J., 1972. Major heavy mineral assemblages and heavy mineral provinces of the central California Coast region. Geological Society of America Bulletin 83 (7), 2099–2104.

Yang, Jingsui & Wu, Weiwei & Lian, Dongyang & Rui, Huichao. (2021). Peridotites, chromitites and diamonds in ophiolites. *Nature Reviews Earth & Environment*. 2.

Zekkos, Dimitrios & Medley, Edmund. (2011). Geopractitioner approaches to working with antisocial mélanges. Mélanges: Processes of Formation and Societal Significance, Geological Society of America Special Paper 480. 480. 261-277.

Zhao, Zheng-Xiang et al., "Review of global 2.1–1.8 Ga orogens: implications for a pre-Rodinia supercontinent," *Precambrian Research* 122, no. 1-4 (2003): 85-114.